

Ret'd by
Olga Hartman
July 5-66 arr. 11

Please return ~~the~~
when you've had
a go at this W.

U. S. A. R. P.

This was typed
for me up at
USARP (NSF)
from my long
hand rept.
submitted to
NSF.

Biological Investigations

Palmer Peninsula

and

South Shetland Islands

1962 - 1963

Waldo L. Schmitt
Research Associate
Smithsonian Institution
National Museum

INTRODUCTION

The following is a resume of the biological investigations carried out at the sites visited by your survey party in and about the Palmer Peninsula, Antarctica, and the South Shetland Islands. These are set forth in 18 sections, I - XVIII, more or less chronologically and geographically arranged under the heading "Site Investigations," as follows:

- I. January 18-20; Marguerite Bay area - Adelaide Island, Base T, Avian Island, Rothera Point.
- II. January 22-25, and March 1; Arthur Harbor area - British Base N, Bonaparte Point and vicinity, Janus Island, Torgeson Island.
- III. January 26-27, and March 1; Port Lockroy - Dorian Bay area - Base A.
- IV. January 27-31; Argentine Islands area - Lemaire Channel, Danneborg Islands, Berthelot Islets (British Base F is Galindez).
- V. February 1-3, and March 2; Paradise Harbor area - including Chilean, Gabriel Gonzales Videla Base, and the Argentine, Almirante Brown Station.
- VI. February 4, 5; Danco-Couperville Island area.
- VII. February 6, 7; Melchior Islands area.
- VIII. February 8-10; Wilhelmina Bay - Svend Foyn Harbor - Salvensen Cove area.
- IX. February 10, 11, 23 and 24; Brialmont Cove, Alcock Island, Spring Point.

- X. February 12, 13; Deception Island.
- XI. February 13; Yankee Harbor.
- XII. February 13; Hope Bay.
- XIII. February 15, 17-19; Seymour Island - Snow Hill area.
- XIV. February 20, 21; Welchness, Dundee Island.
- XV. February 25; False Bay, Livingston Island.
- XVI. February 25, 26; Discovery Bay - Ash Point, Greenwich Island
(for Yankee Harbor, this Island, see section XI, February 13).
- XVII. February 28, and March 3, 4; Ardley Island area, and Potters Cove, King George Island.
- XVIII. March 3, 5; Admiralty Bay.

Weddell
Sea

- - - - -

All listings are of a very general nature. It was not possible to enumerate or identify all the seals and birds in any one area or the animals taken in the dredge or fish traps, the insects or the species of cryptogamic plant life.

The dredged material by force of circumstances, want of time and necessary equipment for handling it, was not worked over as thoroughly as I would have liked. What has been saved in the way of specimens collected by dredge, tow net and otherwise must needs be identified by specialists conversant with the various groups of organisms before their geographic distribution, their relation to their environment and their associates can be discussed, and before our findings can be compared with those published by previous expeditions which worked over or collected in the same areas. A list of tow net hauls made is appended to this report.

Nevertheless, as annotated, it is believed that these listings will

convey some idea of the nature of the flora and fauna, terrestrial and aquatic, at, or in the vicinity of the sites investigated.

The comments, recommendations, and conclusions which follow are my own, based on personal observation, and on discussions with other members of you/ survey party.

IN CONCLUSION

Though terrestrial animal and plant life was more abundant in some areas than in others, no truly barren area was found in the sea.

Seals and their numbers are mentioned when they come to our attention. Some effort was made to note bird rookeries seen by various members of our party and by ship's personnel, and where possible, which was not as often as we would have wished, to identify the species inhabiting them.

Noteworthy bird colonies, mostly penguins, have been noted in the area sections of this report (all but sections VII, XIII, XIV, XVII, XVIII, in which poor visibility, insufficient time for searching, or lateness of season may have kept us from discovering any rookeries of consequence).

Each and every area visited in the course of this survey holds great promise for the marine biologist working on, or with, particular animals or groups of them. Representatives of most, if not all, phyle of marine animals, even though we may not have turned them up in our samplings, are sure to be found in the regions explored by us.

Ice conditions more than anything else will restrict, at times, the operations of those who may follow us, as such conditions hindered us on several occasions. But if we wish to know how antarctic and sub-antarctic organisms live, and where, and under what conditions, and above all know

their kinds and distribution, we must not fear ice, or seek wholly to avoid it. Indeed, to get a true picture, and the whole story concerning any and all of them, we need to go where ice is, though it may add to the difficulties of planned or to be planned operations. The word "hazards" might have been used along with difficulties, but hazards will always be minimal if one goes with proper and adequate equipment and possessed with, or accompanied by, someone with "know-how" born of experience. Do not be too fearful of "ice" in choosing the site of your station or your studies.*

* Don Squires of the U. S. National Museum, who is engaged in working up the corals of "The Ross Sea Fauna" for the New Zealand Oceanographic Institute, is strongly of the same opinion. So far as it is humanly possible we should "pursue" our antarctic organisms through the winter as well as during the few summer months. The occasional open water that occurs in Arthur Harbor and perhaps more open in Paradise Harbor in the winter months, could well render such a pursuit rewarding.

Personally, I favor locating the station as close to the mainland of Palmerland as possible and practicable (this was to be a Palmerland station, was it not?) and not in the South Shetlands. Leave the work to be done in those islands to the British who have already done so much, and published so much on that area.

I also favor sites or areas from which can be reached with greatest facility, and within reasonable distance on foot or by boat, the widest range of types of environment, and terrain, and the greatest variety and

numbers of forms of animal and plant life.

With the foregoing in mind I definitely favor placing the biological station or stations, if there is to be more than one, or a main base with one or more satellite bases, one or all, on a line running from Arthur Harbor through Port Lockroy to Paradise Harbor.

Arthur Harbor as a site for a base or major station comes first to mind as the best possible site on most counts. It is a very favored area, weather, ice, boat shelter, water supply, and perhaps "piedmont" for aircraft landings, abundance, though not necessarily variety of terrestrial life (birds primarily), marine life, and ready access to a great variety of environments within not unreasonable distance. This ready access to a great variety of environments holds good for the other two sites on our projected line of stations, Port Lockroy and Paradise Harbor.

Twenty miles to the south of this "base line" is the Lemaire Channel. Here we spent the night of January 27 drifting about in this scenically breath-taking area where the mountains are as ^{impressive} ~~impressible~~ and at the same time as fantastic or as beautiful in shape and form as any iceberg of imposing size and beauty so far encountered on this journey along the antarctic coast from McMurdo to Marguerite Bay and Adelaide Island to Arthur Harbor and now to this outstanding natural beauty spot.

Enhancing many fold the sheer beauty of our surroundings were the crystal clear reflections in the placid 200 fathoms deep waters of the channel of those majestic mountains rising from the water's edge to snow-topped peaks,

doffing from time to time their caps of clouds, thus granting the "by-stander" an opportunity to photograph them full stature.*

* "in all their imposing grandeur" I wanted to add but dared not. I beg to be forgiven for ~~r~~unning in this "rave" but you would have been moved to feel it, if not say much the same had you been with us that lovely night of January 27, 1963.

Thirty to 35 miles to the south you come to Cape Tuxen and Green Island in the Berthelots with its "...luxuriant growth of moss nearly four acres in extent...by far the largest unbroken (six) patch of vegetation yet found in Antarctica." On the way, having passed Pleneau and Petermann Island, with their Adelie and Gentoo rookeries, shag and tern colonies, and the Argentine Islands, where on Galindez the British Base in this area is located, you also would have seen fields of ice on which numerous seals were lying about.

Eight miles from the western end of the base-line is Cape Monaco (not seen) with its reported garland of six islets, reported as crowded with penguins in season, and 42 miles out of Bismarck Strait you come to Victor Hugo Island, and beyond that the open sea.

Some 50 miles to the north via the Neumayer Channel, an interesting stretch itself, are the Melchior Islands off the north coast of Anvers Island with its inhospitable embayments. In the Melchiors, on Lambda Island, is an unoccupied Argentine Base. Though there was not much on land to interest us, the fishing was good and the dredging promising.

Northeastward by way of the Errera Channel one passes Danco Island and Couverville, and 45 miles from Port Lockroy, Cape Spigot with perhaps the second best (second largest) Chinstrap rookery (next to Alcock) in all Antarctica--and five miles beyond Cape Anna with its reported bird colonies, not seen and not identified as to inhabitants because of poor visibility.

Within 50 miles of the base line, or within a like radius from each of the recommended sites, one can have everything in the way of environment, animal and plant life, that can be had in Palmerland, all that any biologist might crave for study, terrestrial or marine, in or out of the laboratory.

One hundred and 10 miles away to the northeastward also is Brialmont Cove and Alcock Island upon which is perhaps the greatest penguin colony in all Antarctica--all Chinstraps, which I hope will go unmolested for all time to come. He who has to study Chinstraps can do so on Deception Island where there are several hundred thousand, on Cape Spigot (Nunatak Negro) where even more have their home, or in the Paradise Harbor area.

Paradise Harbor I rate as desirable for a land based station next to Arthur Harbor, and Port Lockroy third. Paradise Harbor has much to recommend it, a good site though at present occupied by an inactive Argentine station, "Almirante Brown," proximity to rookeries of three different species of penguins (so reported) though we only came upon two of them in our few days in this area, good fishing and a promise of richer dredging returns than we had from the two samplings we had. Here, perhaps more than elsewhere, it may be possible to study or at least

investigate the life and behavior, distribution, condition and survival of the marine inhabitants of the harbor , pelagic, littoral and benthal, as the tidal and other currents often, if not throughout the winter, maintain enough open water channels to permit boat operations. An attempt in a suitable vessel should be made to prove or disprove this contention.

Port Lockroy deserves better than a third place rating. It hurts to put so lovely a place so low, especially in view of its swell-free harbor in which a vessel might safely be wintered (shall have to ask Capt. Mac about this), and where we found the littoral faunal so very rich; but Paradise Harbor I feel quite strongly has the far greater biological potential.

Within Arthur Harbor, I favor the British Base, that area or site over Bonaparte Point, or the so-called Bravo site, for reasons stated elsewhere in this report, particularly in the discussion of the Arthur Harbor (section II, pages 3, 4).

I would urge that we endeavor to get full and clear "title" to all three places by transfer, as gifts, or by some reimbursement, permitting the present builders, or owners of the huts in sites in Port Lockroy and Paradise Harbor, to remove buildings and contents should they care to do so. We should by all means start anew, and with our own new buildings or structures.

And now? What are we waiting for? Let's go!

POSTSCRIPTUM

Last but not least - ever since I visited McMurdo Station - I have become enamoured with the idea of a floating laboratory, or at least a sufficiently large trawler-type of vessel to carry on the lines of work planned of any shore-based laboratory that might be set up in Palmerland. These notes I shall organize at the first opportunity which has not yet (the time to do it) come to hand - give me just a little more time. This survey is but one of several very worthwhile projects that I have on the "fire" at this writing!

Appended to this report are a list of the two-net hauls or tows made in the course of the survey; and an outline itinerary for ready reference. Appendixes "A" and "B".

SITE

INVESTIGATIONS

Marguerite Bay area - Adelaide Island, Base T, Avian Island, Rothera Point

The ice fields through which we passed getting here had quite a number of seals on ice cakes here and there. Mostly, if not all, Weddells, I take it. Some said they saw a leopard seal, but I saw none that quite fitted the description given.

Seals must be plentiful in this area; it was said that about 500 a year are killed for dog food and station mess.

At all places the "usual" birds, many still nesting here: Adelie penguins, skuas, blackbacked (Dominican) gulls, blue-eyed shags, and a giant petrel; at Rothera Point also, antarctic terns.

Mosses and lichens were collected at each place; from them mites and spring-tails were Berlesed; got about three different species of insects.

From a fresh-water pool on Avian Island a few fairy shrimp (Branchinecta) were dipped with an empty beer can lying about (Tom Berg, coll.).

Fresh-water and marine algae were collected; and at Rothera Point red and green snow algae as well.

From fish trap set in five fathoms off Avian Island, no fish, a number of nemertean worms, amphipods, 2 species of starfish; 3 small sea urchins, these last at Rothera Point where the worms and amphipods also appeared in the traps.

Dead limpet shells left by the gulls whose principal food are limpets seemed to be scattered about everywhere, some with serpulid worm shells attached.

There was much ice floating about, making tow-netting difficult. This was said to be the best year in the last six, more open water than usual. However, the ship had to shift position to avoid a sizeable iceberg; can be a nuisance here.

Avian Island despite its teeming bird life, fresh-water pools, many mosses, is not to be recommended as a station site because of the interference with the penguin rookery (and the "mess" it would have to be located in; no suitable space not occupied, but I did not get over much of the island).

Base T site hilly; good buildings, best is the one-man ("weather shack," ideal set-up of what a one-man study and laboratory could be; and sleep-in.

Rothera Point, though, has the building land (acreage and "foundation-wise," a fine place)--a low, level saddle between the "hills" toward the Point and the more mountainous piedmont to the right as viewed from the Bay landing. I wonder how wind-swept this place is; looks like a "draw." We had a fine day here, only a windy and stormy one would tell.

There were only a few penguins here compared with the hundreds at Avian Island. Lt. Thomas attempted to walk around the shore of the Point but gave it up because of the determined way the skuas, which must still have been nesting, dived at him; although he saw evidence of what might have been a penguin rookery but no birds on it. Quite a few small bergs about in the landing area; seals on a number of them.

January 22-25
March 1, 1963

Arthur Harbor area - British Base N, Bonaparte Point and vicinity, Janus and Torgeson Islands

There are quite a number of seals about; we had no difficulty in locating a Weddell on an ice cake for bait.

The area is rich in bird life, especially Adelies. There is a large and populous rookery on Torgeson Island, and other sizeable colonies scattered through the area. About Cap Monaco to the west are six islets with rookeries on them; there is also a colony at Biscoe Bay on the S. W. coast of Anvers Island (fide sailing directions); others on Litonfield, Halfway and Humble Islands (fide Berg helo flight), Giant petrels, skuas, Dominican gulls, as always and in considerable numbers, Antarctic terns, and sheath-bills. Many dead limpet shells ashore, others gathered from rocks at tide level.

Mosses and lichens, grass clumps collected, and insects berlesed from this vegetation.

Fresh water algae and fairy shrimp were found in the pools atop the bluff above the Base N hut. Along shore at Bonaparte Pt. and at Base N marine algae and a few amphipods were dip-netted.

#7-63

Dredge hauls were especially productive - a varied and abundant fauna on a stiff blue mud bottom, 12-17 fathoms, - a species of sponge, sea urchins, 2 sp. starfish, worms in number, the nemerteans proved to be the common species throughout the area (Amphiporus), 2-3 species of annelids (Nereis, Terebella and at least one other), 4 species of mollusks, and

several species of amphipods, an isopod, and 4 or 5 species of sea squirts (ascidians).

#67-63

The dredge haul of March 1, on our return to Arthur Harbor was a repeat of the earlier one with a host of annelid worms and mollusks.

#8-63

The fish trapping was equally rewarding, 88 notothoniid fish were taken in two sets off Janus Island together with starfish and nemertean worms. The fish ranged from 7 to $14\frac{1}{2}$ inches in length. One trap set in the rocky part of Bonaparte Inlet was crushed under a small berg that either drifted in with the tide or possibly was calved off during the night from the ice cliffs lining one side of the head of the inlet; considerable sea weed and a large clump of ascidians were dislodged as we got the jammed trap up after some hours' labor, dried in forenoon, were successful with a grapnel in the afternoon.

Brash-ice troubled our March 1 landing at the Base N site, recent snows had whitened the ground and the large lake a few hundred yards behind the hut was thin-ice covered, as were the fresh-water pools on the top of the bluff above the hut. A few lonesome-looking Adelies were standing on the sea-ward side of the bluff; skuas were about but in lesser numbers than in January. Where do they go; migrate as do the penguins? Torgeson Island was all but deserted, just a scattering of penguins about.

The approaches to Arthur Harbor, because of the current running through it, is said to be quite free of sea-ice more or less the year round.

Regrettably during the January two-hour squall we experienced in Arthur Harbor, we did not get to see the Inlet; it is a remarkably protected,

quiet bit of water. On March 1st like a millpond, though it was windy and rough out in the harbor.

I do not particularly fancy the Bonaparte Point site because of its exceeding rockiness - large jagged, broken rocks, and a rather uneven gullied (or is it ridged) terrain. The Inlet, though a piece of very quiet water beautifully protected from wind and wave, might be a very dangerous place. It narrows down to about 80 feet in width, and on the right hand side are ice cliffs of about that height. These have plenty of fissures; cracks, and crevasses. A fall of a mass of ice and snow could well spell disaster to any vessel that might find itself in the Inlet at the "right" (or would it be the wrong) time. Really, how very accessible is the piedmont here?

Getting fresh water would be more of a problem than at Base N where it could be conveniently piped from the large and quite deep fresh water lake there. No dam would be needed, the valley in which it lies could hold many times the present amount of water without damming; or much of one, but the supply of water seems ample for the needs of any station that might be established here.

Within walking distance and not near enough to be objectionable, to one side is a penguin rookery, on the other the bluff on top of which are situated the fresh water pools already referred to. A man needs to get about a bit for relaxation if for no other reason. At Bonaparte Point he could walk about on rocks and still not "be" or get anywhere.

True, the boat landing is not of the best but it could as easily be

improved as that which would have to be provided at the Bonaparte Point site.

During the January squall the Navy Survey Party was caught out at the Base N landing. Lt. Nash said it was "not so bad" and that there was some degree of protection from what went on in the harbor outside.

I might add here, too, that Lt. Nash also favors the Base N hut site over Bonaparte Point from an engineering point of view. I have his permission to so state.

On 21 January we were underway for Anvers Island and Arthur Harbor.

January 26-27
March 1, 1963

Port Lockroy, Dorian Bay area, Base A.

In this area the Gentoo penguins take the place of the Adelies but for this major difference bird life is as varied as elsewhere, although the number of individuals seems less than at Arthur Harbor. Three chin-straps were seen among the Gentoos here by Lt. Thomas. I do not recall seeing any giant petrels here. Rookery space is more limited, resulting in fewer colonies, but those available are well populated. Near at hand to, but not on the Base A site, are sizeable Gentoo colonies, and also at Damoy Point on the way to Dorian Bay is another large one. A colony of terns holds forth on a rocky, steep-to headland near the entrance to this port, and a well populated shag colony occupies an island in Peltier Channel (Priest Id?) on the way to South Bay, Doumer Island, where the Chilean Base "Yelcho" is located. Additional colonies of Gentoos occur down this way, also. Skuas, Wilson's storm petrels and sheath-bills were about, too, in the harbor.

There was comparatively little vegetation to be seen, nothing at all to compare with the luxuriant vegetation that we were to see later in the Argentine - Berthelot Island area. From moss gathered here a few Collem-bola were Berlesed.

#9-63
#66-63 The more meager dredge hauls here, over a muddy bottom, in general character resemble those made at Arthur Harbor. Worm tubes were abundant; Ophiurans (serpent or brittle stars) and ascidians of species other than those obtained at Port Arthur were washed from the mud. Additional specimens

of these species were secured by the same means on occasion of our March 1 visit and, unexpectedly, a tiny gasteropod.

#10-63

The richness of the shore fauna is worthy of special mention. It so happened that we landed at the Base pier at extreme low tide. In the course of a few scrapings along the rocky shore Poblete's dip-net brought up a variety of animals - the already (to us) common nemerteans (Amphiporus) three sea-anemones resembling the North temperate Metridium (the only sea-anemones of the trip, it turned out), two species of amphipods, and two widely different and very characteristic antarctic isopods, Serolis and the giant, almost 10 cm. long Glyptonotus antarcticus. And had we had as favorable a tide in the Arthur Harbor area, I like to think that we would have been as fortunate there.

#11-63

From the fish traps we got more nemertean worms than we knew what to do with, 2 species again of amphipods, and 7 notothoniid fishes.

The building area about Base A (Goudier Island) is a rocky one but nowhere as rough going as on Bonaparte Point, Arthur Harbor. Port Lockroy is one of the few ~~swell~~-free Palmerland harbors. As to aircraft facilities, landings, see Capt. McDonald.

I cannot refrain from adding that Port Lockroy is as scintillatingly beautiful a place as I could spend an antarctic summer in, or a winter, too, for that matter - Lemaire Channel is another.

January 27-31, 1963

IV

Argentine Islands area - Lemaire Channel, Danneburg Islands, Berthelot Islets (British Base is on Galindez)

Lemaire Channel, Jan. 27-28; Pleneau Island, Jan. 28; Petermann Island, Jan. 28; Galindez Island, Jan. 29; Cape Tuxen and Green Island, northernmost of the Berthelots, Jan. 31.

Galindez Base gets its fresh water from a large pond (lake?) behind the station. Many seals were basking on the ice cakes, thick in all directions, south of Galindez, more seals over a larger area than any in which we had seen any large number of them. Two whales were seen north of Galindez on the 29th.

An interesting and well populated area, animal, as well as "vegetable"-wise. Many birds breed in this area. The following list was given to me by Mr. R. Lewis of the British Base here on Galindez:

Dominican (black-backed gull) - a year-round resident, an interesting record; great skua (breeds); antarctic (or is it the swallow-tailed?) tern (breeds); giant petrel (visitor only), always seems to appear in bad weather, several to half a dozen of them; sheath-bill (winter visitor); Wilson's storm petrel (breeds); snow petrel (breeds locally); Adelie penguins (are on Jalour and Petermann Islands); Gentoos also on the last-named island; chinstraps (casual visitors to the area); Emperor Penguin (seen once); blue-eyed cormorant (shag) (breeds). A cormorant colony, not seen by us, is located on Winter Island.

On Peterman Island both Gentoos and Adelies have rookeries; among the Gentoos one Adelie made itself at home. On a rocky peak across a snow field

behind the Gentoo rookery, antarctic terns were flying about in considerable numbers, some were nesting. Jack Crowell saw one with a chick (close up, - beauty of this nesting site was the fact that you could walk onto and around it to photograph the birds close-up).

At Cape Tuxen there is a colony of blue-eyed cormorants.

On our return to the ship from Green Island we tried to get photographs of a flock of 30-40 (? 20-30) of these shags sitting on the water quietly and in a surprisingly close group, but they proved wary; before we got into comfortable camera range, they dived and scattered. Green Island has a considerable population of skuas. Judging from the aggressiveness displayed by several of them, they were protecting nests and young. A large colony of cormorants is also located four miles south of Cape Tuxen in the Berthelot Islands.

Mosses and lichens were gathered from the several islands and the Cape, and Berlesed. An amazing number of Collembola were driven out of a couple of handfuls of moss from Green Island (I brought this vial along to show you).

At Cape Tuxen and more so on Green Island, the moss growth was especially luxuriant. Most of the islands, to a greater or lesser extent, are "green as grass" on their northern, more or less snow-free, slopes.

Green Island, the northernmost of the Berthelots, lives up to its advance notices in the Admiralty's Antarctic Pilot (p. 204, 2nd edit., 1948), "....Green islet has on its northern slopes a luxuriant growth of moss nearly four acres in extent with peat up to three feet (o m g) in thickness. This is by far the largest unbroken patch of vegetation yet found in Antarctica.

The "peat," so-called, was thick, but none that I saw or walked over was quite three feet thick, nor was the moss so unbroken. Where the moss grew over and down the sides of a boulder or heap of rocks, it might easily give one the impression of being a moss clump or heap three feet thick. It was thick, all right, much of it was dead or looked so underneath. There were breaks and gaps in the patch as a whole, but there were no extensive gaps or large areas on the slope bare of moss - quite a sight!

#14-63

Two dredge hauls made at our Galindez Island anchorage, in 25 to 42 fathoms, and results combined into one. They brought up a portion of a silicious sponge, a number of worms, starfish ophiurans, a red sea-urchin and the first crinoid of the cruise, a gastropod mollusk, and a number of long-stalked ascidians. A second specimen of the same species of crinoid was found in mud attached to the anchor when it was raised on 31 January.

The fish traps disappointed us, no fish, nemertean worms, a few, and a starfish,--the cove by the Base landing was evidently a poor location for catching fish. Had there been time a second set in another location might have brought us better luck. We had drawn a blank as regards fish in Arthur Harbor the first time around. It was the second at Arthur Harbor that turned up the best catch of the cruise.

This area with its many islands and its nearby "deeps" and channels (Lemaire) is undoubtedly a very rich one biologically. For a possible satellite station in this area a site with a good boat shelter should be selected, Winter Island - Stella Creek where the Penola wintered, Petermann Island where Charcot holed up in 1902, or possibly Cape Tuxen at which another look might be had for faunal and floral growths - it is

hard-by Green Island. But then again, I do think we should leave the area largely or wholly to the British; they were here first, have done a lot, and will undoubtedly do more as time goes on. Not long before our arrival, the folks at the Base had sent off quite a lot of "pickled" invertebrates to the British Museum. Here should be recorded also, that to help me out with my shortage of bottles they gave me a case of "handsome" wide-mouth gallon jars - a "God-send," if ever a biologist had one!

February 1-3
March 2, 1963

Paradise Harbor area - including the Chilean Gabriel Gonzalez Videla Base and the Argentine Almirante Brown Station.

This area is said to have breeding colonies of three species of penguins. I have no reason to doubt this but I saw only two, Gentoo and Chinstrap, in goodly numbers and in several places. Members of the Chilean Base said the Addies are not far away at "Paterna," which was written out for me. This locality, if it is one, I have not been able to locate. I expect by correspondence later to run it down. There are extensive shag (blue-eyed cormorant) colonies in this area or nearby on Cape Duthiers, "just around the corner" on the way to Anvord Bay. Cape Pigeons, and about all the birds seen elsewhere.

The Chilean station is built upon an island thickly populated by Gentoo and Chinstrap penguins, right in among them. Seeing what's here, what it has done to the birds, in part at least, and the filth that surrounds the buildings as a result, I would recommend against locating any station on a bird rookery; it is neither fair to man or beast.

On a point across the harbor on Lemaire Island is a penguin rookery (birds not identified - did not get close enough), and on Bryde Island on a rocky ledge or islet hard by is almost a good-sized penguin rookery, around the shores of which a number of seals that I take to be crab-eaters, had hauled out.

Mosses collected yielded Collembola in modest quantity.

Two dredge hauls made on rocky bottom, 33 and 41 fms., returned a variety of invertebrates, perhaps the greatest variety seen so far; sponges,

#17-63
#24-63

hydroids, and another type of coelenterate polyp that I cannot at the moment readily place, bryozoans (moss animals), ophiurans, holothurians, clams, two caprellid amphipods - elusive creatures - the only two recovered from any of our dredgings, and ascidians again. Palmerland waters, if not the rest of Antarctic, is certainly sea-squirt heaven. I have collected many over the years but never so many different kinds, so structurally different (external appearance) on any previous cruise.

#20-63
#25-63

Of two sets of the fish traps, the first off the Chilean Base, we got nothing but a "mess," a 100 or more nemertean worms and several different species of amphipods. A large jelly fish and a ctenophore, "comb-jelly," were picked up drifting by in the tidal current.

With the dip net a lot of algae and amphipods, a large red species (Paradoxmine ?) were captured. These could well furnish ample meals for many penguins. I get the idea from the penguin behavior here. As we were working along this stretch of shore several penguins were seen actively diving, duck fashion, heads down, tail up, and when their heads came up above water again, working their bills as though they were eating something that either tasted good or was satisfying. After a number of such dips, these birds, three or four of them, headed out to sea. I did want to have a look into these particular penguin stomachs to verify my suspicion that this omnipresent and abundant food material formed an important part of the penguin dietary. If the amphipods are as abundant elsewhere as in the red sea-weeds here, they must play a far more important role in the economy of penguin life than heretofore realized (or has this side of the penguin dietary already been studied by someone else?).

As far as our tow-netting efforts are concerned "krill," the Euphausian "shrimp," have been elusive animals, but as luck would have it a goodly number were thrown up on the forecastle deck in the small hours of the morning on our way to Paradise Harbor; one was still alive and swimming about in the depression under a flush hatch-handle. From the anchor the same morning a small, white holothurian.

I am much taken with Paradise Harbor and the opportunities that it may have for biological research. I feel sure that biologically at least it will justify its name.

The harbor has its ice conditions but I read in the coast pilot, or sailing directions, that although there is considerable movement of ice through the passage between Lemaire and Bryde Islands, the upwards of 3-knot tidal currents prevent ice in the bay from freezing solid in the winter months (could be). Bergs there are, and more will be calved, but do not let us forego the studies that should be made of animals under natural conditions just because of some floating ice. We should have the whole year-round picture so far as possible, not just a series of summer sketches.

Moreover, Paradise Harbor indents the mainland, not one of its many off-lying islands.

There is an excellent station site, that of the Argentine Almirante Brown Base, with not unworthy boat shelter nearby. Hope we can get title to the place and erect entirely new buildings. Against the high cliff face, I should like to see a multi-story building, two or three, if not more, stories, with a cantilever platform "out front" to which the station boat could be hoisted, as well as supplies that could be stored right in the first story of the building.

Such a building would be as snow-free as the cliff to which I would attach or anchor it. All problems, waste and water, would be solved (snow and ice for melting are atop the cliff or close by), plenty of ice and bergy bits float by. The several storied building would make for the conservation of much of the warmth (heat) lost through the roof of a one-story building. Then again, the fact that the cliff side of the building is not exposed to the winds would also minimize heat loss.

Danco - Couverville Island area.

Here is an outstanding bird area - Gentoo and Chinstrap penguins in large rookeries. The former are centered on Couverville and thus negate this excellent building site as a place for a laboratory. There also happens to be a fine safe boat shelter on the east side of this island behind a natural breakwater upon which a small shelter could be erected. At high tide it would be isolated as water flows in over the in-shore end of this ridge. Atop its widest part is an abandoned whale boat. The whalers knew a good shelter when they saw one. There is quite a bit of snow-free rock here. Jack Crowell has walked over it.

A short distance to the north of the Base "O" hut, on Danco Island, there is a large Gentoo colony, where birds were marching back and forth in deep "ruts" or tracks all the while we were ashore. The landing is poor here because along the shore the water is so very shallow for quite a distance out.

Last but not least is the Chinstrap penguin rookery, or rather metropolis, on Cape Spigot. It is a sight worth travelling to Antarctica to see. On the Argentine charts the name is Nunatak Negro. Its foreshore is crowded with Chinstraps, among whom a single Gentoo was spotted, but the bulk of the Chinstraps in this vast rookery had nests, or roosts if you will, all over the rock, steep slopes of Cape Spigot. Every snow-free patch of rock, and the greater part of this great and impressive Cape was crowded with penguins. Capt. McDonald estimated that this Nunatak must have at least 300,000 Chinstraps. What hikers and climbers the Chinstraps must be to avail

themselves of this high and steep-sided peak to colonize its uttermost heights, 938 feet above the sea! Most of the road upwards for those that had not settled on the foreshore was over a small steep-to snow field lying shoreward of the bare crest of the "saddle" between the peak of the Cape and the ice sheet, or cap, further to the north of it.

#27-63
#28-63

Two dredge hauls were made on February 5, the first at the Base O, Danco Island, in 41 fms.; the second at the anchorage in Errera Channel nearer Couverville in 46 fms. At both the bottom was more or less rocky and very much alike in their sampling of the channel bottom animal life; Echinoderms predominated, many ophiurans, a few starfish, and a dozen or more red sea-urchins. In the second haul were several forms of animal life not found in the first, a large nereid worm, a number of bryozoan fragments, and what made me want to let out a cheer, a hippolytid shrimp, the first and, as it proved to be later, the only decapod crustacean taken on this cruise. Two white starfish came up on the anchor at the first of our anchorages of this day. A small silver-sided fish, also unique in our collections, 6 inches long, apparently dropped by some bird, was picked up on Couverville by Lt. Nash while ashore surveying a possible site. No really suitable site for a station of any size seems available either here or at Cape Spigot unless one wants to bed down with the penguins. The possibility of finding space enough for some small shelter on the natural breakwater on the east side of Couverville was mentioned above. The only need would be for penguin studies, the marine fauna would be a job for a vessel hereabouts, and Chintraps might be more conveniently studied in Paradise Harbor if and when one might acquire the Argentine Base there; and as we learned later, at Deception Islands.

Ink notes added 8.VI.66

VII

February 6, 7, 1963

Melchior Islands area, principally with reference to the Argentine Station located here on Lambda Island.

There was a great deal of snow about and between some of the buildings - most of the other islands round-about and this one off the station area are heavily ice-capped. Not much chance to set up "business" here or the urge to do so after seeing this site.

see note bottom of page, ~~me~~ but
I do have note that ~~some~~ things were
picked up

No shore collecting done, no real opportunity for it while here, few birds about, some Dominican gulls, Wilson's storm petrels, and a giant petrel (brown phase).

4-32-63

Our dredge, hauled over a mud and sand bottom at 25 fms., brought us the first sponge of any appreciable size so far, a compact, siliceous one about 8 or 9 inches in diameter; also worms; a starfish; a number of sea-urchins; an alcyonarian; a bright yellow sea-slug; half-a-dozen sea-squirts; and a number of tiny red amphipods that either lived on that siliceous sponge's rough surface or in its canals.

34-63

33-63

From the fish traps, after an overnight soak in 7-11 fms., 22 fish ranging from 11 to 14½ inches were found, together with two starfish, and a "crop" of the omnipresent amphipods. Four other fish were taken over the ship's side on hand lines. On the anchor, as it was hoisted, were two bright yellow prickly (soft-spined) ascidians like the bright yellow sea slug in color, also the first of their kind seen on this trip. Is there any reason for this identical color in two widely different forms of animal life - from the same general area, and at that not so far apart?

His
is
str.
33

(X) Lambda was the only island visited during the short time spent in

this area. From a marine biological point of view the biological potential

at bottom of note book page I have "~~couple~~" ^{on which notes for 32-63 were entered} entry:
 "# 32 a amphipods and small stuff from shore."
 This must have been shore of Lambda Id.

is high and holds promise of even more interesting things to come than bright lemon yellow invertebrates representing two widely different phyla.

On the way from Melchior to Port Lockroy the ship headed southeastward down the Schollaert Channel. On the way, she entered each of the great bays indenting the northern coast of Anvers Island; first Lapeyrere lying between the Hump and the Gourdon Peninsula; next Patagonia Bay between the Gourdon and Thompson peninsulas; and lastly, Fournier Bay and Inverleith Harbor which indents its eastern shore.

Some seals, not many in any case, were basking on ice cakes or small bergs in each of these bodies of water; a killer whale (Orca) was seen in Patagonia Bay - little else, only a few stray birds, other than the omnipresent Wilson's storm petrels, a Dominican gull or two, a skua now and then, a tern, a giant petrel, and a snow petrel or sheath-bill. Saw more seals than birds. A foggy overcast, somewhat dismal day; ice cliffs round about.

Spotted no "sites" likely or unlikely.

Wilhelmina Bay - Svend Foyn Harbor - Salvensen Cove area.

In the afternoon of the eighth, the ship moved into Wilhelmina Bay past Cape Anna. The Cape is reported to be the nesting site of countless cormorants and gulls but distance and not the best visibility precluded identification of the birds whose rookeries we glimpsed.

Capt. McDonald, who flew a helicopter reconnaissance of Wilhelmina Bay while the ship was lying-to, reported: Ice cliffs on all sides, no building sites or boat shelters, two active glaciers, some fast ice.

The night of February 8, the ship anchored in Svend Foyn Harbor at 1940. We then took out our fish traps and made a tow-net haul. The dredge came up empty this evening, so the haul was not counted and no other attempted because of the lateness of the hour.

Shore collecting also proved impractical this evening and the next day when a successful dredge haul was completed and the fish traps lifted.

#37-63

Dragged from 18 to 25 fathoms over sand and gravel bottom. The dredge bucket brought in a foot-long nemertean worm; several annelids; some small white starfish; a considerable number of red sea-urchins of the species we have taken on a number of occasions; a rather large ophiuran differing, so far as I could make out, from all previously caught; a number of bryozoan fragments; the first brachiopods of the cruise, quite tiny fellows though; two species of mollusks; a few amphipods (as usual); and about a dozen sea-squirts.

#36-63

Thirteen notothoniid fish were trapped. Of these the largest measured

15 inches in length, the smallest, of apparently another species, $5\frac{1}{4}$ inches. A 10-inch notothoniid was caught over the ship's side on a hand line.

February 9, Cmdr. Lewis gave me a photo-copy of this area, on which he had noted the penguin colonies he had seen in the course of two days of helicopter reconnaissance flying; he also noted any conspicuous lichen stands that caught his eye -- encrusting yellow lichens primarily. The flights ranged from Hansen Island northward as far as Challenger Island just above Bluff Island. His observations, as entered on the charts herewith, follow: He considered as "small" colonies estimated to contain under 1,000 birds, and as "medium" if the individuals present were estimated to number from 1,000 to 5,000.

- (1) North end of Delaite Island, small penguin colony.
- (2) Two rocks, or islets, south of the largest of the Racovitza Islands, have each small penguin colonies on them.
- (3) (On the way into the cove in Svend Foyn Harbor in which the wreck lies, a rocky point to the left has a shag colony on it--half the birds were sitting, perhaps better standing, in the sheet of snow above the bare rock area on which the rest of the birds stood.) (This is my personal observation entered on the Commodore's charts.)
- (4) Penguins on rocks and islets ringing Icarus Point.
- (5) And at the western end of Bancroft Bay numerous shag colonies.
- (6) Left "arm" of Reclus Peninsula on a 500-foot cliff (west of the 1270 elevation on the chart) a medium-sized penguin colony.
- (7) Southernmost of the Gaston Islands medium-sized penguin colony.
- (8) Shag and sparse lichen growth on Andree Island in Recess Cove.

(9) On Peninsular extending out from shore before Santos Peak, a small penguin colony either side.

(10) Northern extremity of Bluff Island small penguin colony, and lichens.

The Commodore also indicated that there were many seals on, or about, the rocks (and/or ice) and islets north of the Reclus Peninsula, and about the little hooked peninsula, Portal Point, extending eastward, and more seals a little farther south along the coast.

Sven Foyn Harbor was much favored by whalers in by-gone days. It is said to be a place of generally favorable weather (H. O. Sailing Directions, p. 194), even though there are at times violent southeasterly winds. After having seen all three places in this area, Capt. McDonald believes that ice conditions are probably better here than at Alcock Island farther up the coast, or at Welchness, Dundee Island, over on the east side of the Palmer Peninsula.

The Captain also remarked that there is a small boat shelter available here near the islet on which rest several of the old "man-powered" whale boats or what is left of them. On this small rocky islet is space that could be utilized for a small building (sub-station or shelter).

The bottom of the harbor is rocky, and poor holding ground, a fact appreciated by those whalers for they installed moorings here and there on rocky ledges as elsewhere in places favored by them along the Palmer coast and islands.

The Staten Island moved out of Svend Foyn at 1500 and cruised the Brabant coast the afternoon of the 9th. No site areas noted. Passed Auguste

Island and Cabalescou Islet. The former is mostly snow-free, the latter, also largely snow-free, has penguins and cormorants "aboard" (fide sailing directions), but we were too far off to see what was what. The former small island is beset with shoals and apparently offers no shelter for small craft.

For the night we lay over in Hughes Bay and the ship's force spent the following forenoon overhauling the helicopters, effecting repairs to one of the two steam boilers, and shifting the LCVP #1's motor into the Greenland cruiser.

As the ship passed through Salvensen Cove one of the deck officers noticed several patches of "red water" but did not mention the matter until some time later, too late to do anything about it. It assuredly must have been krill which has been eluding our nets and drags except for an occasional specimen. How I would like to have made a tow through that "red water."

On the starboard side going into Salvensen Bay there is quite a large cormorant colony with half the birds sitting up on the snow field above the rock exposures that marked the nesting sites. It was like the state of affairs we had earlier observed on some exposed rocks at the head of a cove harboring the wreck in Svend Foyn Harbor.

Dredge haul and fish trap evidence backed up by "red water" seem to indicate that the Svend Foyn area might profitably be exploited biologically.

By 1100 the ship was underway again, headed for Brialmont Cove.

February 10, 11
23 and 24, 1963

Brialmont Cove, Alcock Island, Spring Point.

Several hours were spent locating a suitable and satisfactory anchorage; a considerable part of the cove is around 200 fathoms deep.

This is quite a lively place, seals scattered about. It has been some time since we have seen quite so many around any of our anchorages. They are mostly, or all, crab-eaters. Capt. McDonald said that he saw one leopard seal in the bay, and a number about Alcock Island on which there is a large penguin colony.

We killed one of the crab-eaters for bait for our fish traps. Doty shot him; the very first shot must have severed the spinal cord, for the seal quivered and died. Doty's second shot was not needed but was fired as a precaution before the boat crew climbed over onto the ice cake where the seal was resting so that we could get this "ton," it seemed, of seal meat aboard. In its stomach were several gallons of krill. The krill in the two ends of the stomach were distinctly different in color, and I believe due to a difference in species rather than degree of digestion. Shall check when specimens get home.

Contemplating that seal's relatively small mouth and insignificant teeth, and then the gallons of shrimp in its stomach, one is forced to conclude that the krill were so numerous and so crowded together that the mass of them must have had the consistency of thick porridge. How else could that seal have picked up perhaps three gallons of small shrimp in his small mouth?

Again, the usual run of birds was found in and about Brialmont Cove and more numerous than usual: Storm petrels, Wilson's; Cape pigeons; Dominican gulls; sheath-bills (shags); skuas; a giant petrel; and penguins, and penguins! The latter appear to have taken over about all available rocks and islets about the Cove and virtually the whole of the snow-free parts of Alcock Island. The penguins are Chinstraps, the little fellows with the urge to climb high in this world of ours. Alcock's difficult and precipitous terrain reminds one of Cape Spigot in the Danco-Couvertville Island area, but nowhere nearly so high, only a bit more than 300 feet, as compared with Spigot's 938. But Alcock must be much more densely populated. Capt. McDonald, who has reconnoitered Alcock from the air and has landed on its shores, thinks it has more penguins on it than Cape Hallett, with its 300,000, down McMurdo way. He said he believes it to be about the largest penguin rookery that he has ever seen in all his eight years of antarctic experiences.

At the foot of Alcock's precipitous slopes, the Captain found a small cove that would afford shelter for a small boat. Adjacent was a piece of penguin-free land upon which one might build without interfering with the Chinstrap way of life, which includes an "awful" lot of steep, uphill climbing, for they occupy almost every bit of snow-free rock on Alcock, as at Spigot, to its very top. However, I would like to see the Alcock Chinstraps, as well as those at Cape Spigot, left undisturbed. Anyone having the urge to study Chinstraps can do so in Paradise Harbor, living as did A. C. Bogdanovich at the Chilean Station there in 1959-1960 as an observer, or even more conveniently at Deception Island where live several hundred

thousand Chinstraps, and where comfortable accommodations may be found at any one of three stations, Argentine, British, or Chilean.

Mosses and lichens were collected ashore on each of our visits to the Cove, February 10-11, and 23-24, at Alcock Island and at Spring Point. In spray-fed pools, or rather pockets of water, among the rocks on a rocky islet that is virtually a part of Alcock, we dip-netted with a fine meshed net the young stages of some aphipod, along with the algae that were growing in the same little bodies of water.

#40-63
#58-63

Two dredge hauls were made here. The first on February 11, in 35 fms. off the ship where the bottom may best be described as a "regular concrete mix," sand, gravel, and good-sized stones. Some of the stones and larger pieces of gravel carried encrusting bryozoans. There were also a few small worms, amphipods and bits of algae in the dredge. That our dredge did not bring up more bottom-dwelling organisms may have been due to the inadequacy of our equipment for this type of bottom, or that the place in which dredge was dropped was rather barren. Only another drag might tell. That opportunity was vouchsafed us on the 24th of February. This time there was mud mixed in with the gravel in a depth of 18 fathoms. The dredge when brought up was quite alive with a variety of invertebrates: three or more species of worms; disintegrated bits of some jelly-fish-like organism; hydroids; an aloyonarian or antipatharian; both branching and latticed bryozoans; starfish, one tiny opiuran; and several crinoids, the first time that we have taken them in numbers.

What might we have not gotten in other parts of the cove if there had been time and the facility for carrying our ^{"oil-drum"} dredge about in it.

We almost lost one of our fish traps here beneath a small iceberg. We were a long time getting it out from under, and when we did get it up, it was practically empty. From the two traps set we got only a few amphipods and bits of algae.

Despite the rather poor showing of dredge and trap, I still think that we have an area that will prove rich if ever intensive work is carried on in it.

With a suitable vessel and the right gear, one would not need a shore-based laboratory to properly work over this area. I would recommend the acquisition of such a vessel.

Deception Island.

This place has most everything, plenty of "flat" land for building sites and airstrips, and plenty of running melt-water - even a "built-in" source of heat, and power perhaps. Thermal wells produce heat and power in New Zealand's North Island in Italy. Why not here where clouds of steam arise at low tide from the beach sands down in front of the old whale factory?

For him who wishes to study Chinstraps, this should be the place. Some 250,000 Chinstraps live on the outer slopes of this old volcano. Excellent accommodations could probably be arranged for at any one of the three bases or stations maintained here by as many different nations, Argentina, Chile, and the United Kingdom.

A number of other birds are to be found nesting here as well. Cape pigeons are around most of the year except between August 11 and September 8. Common also is the Dominican gull, which in contrast to its fellows in the Argentine Islands much farther south, is a non-resident at Deception Island. At Deception, we find also sheath-bills (or snow petrels), Wilson's storm petrels, the giant petrel, swallow-tailed (or antarctic terns), and blue-eyed shags. Gentoo penguins are occasional; sometimes "touring parties" of 10 or a dozen arrive, at times accompanied by single Adelies. At times macaroni penguins are seen; three pairs have been seen nesting among the Chinstraps. Bird-wise, this is an altogether interesting place.

In the course of a brief period ashore three or four sheath-bills were noticed picking over a pile of rocks on shore, and behaving along the water's edge like so many waders. Scrapings from the rocks were made. All that we

noticed were some tiny mollusks (gasteropods), and a coating of green algae. Mosses, lichens, and fresh-water algae from a melt-water stream were also gathered in; the mosses Berleazed for insects.

Time was not vouchsafed us for a dredge haul. It is true that with the trash and "junk" on the bottom left from the whaling days, we might have lost our dredge, but made aboard here by the engineer's force out of an old steel oil drum, replacement would not have been difficult, and there was cable to spare on the hydrographic winch, the outer turns of which had been pretty well used already.

Fishing with the traps revealed that fishes were plentiful. Thirty-five were taken in the overnight set in 5 and 8 fathoms. They range in length from 11 to 21 inches; the larger fish were in the trap set in 8 fms.

Near the shore before the old wrecked whale factory and try-works are a number of large steel tanks, rusty to be sure. A couple of them have had openings cut into them at ground level and are used as store houses. Should ever a "station" or laboratory be planned for Deception, it would be an easy matter to convert one or more of the remaining tanks into laboratories and quarters. Cork insulation as we have it aboard the Staten Island, ports and doors cut in, two decks installed, would give one a very substantial, comfortable, already roofed buildings with a possible source of heat hard by the door, and water or its makings close at hand.

There is a well, back from the beach, in which the water has a temperature of 65°F, but this is not for drinking, melt-water streams are everywhere, and snow and ice readily accessible.

Except for the fact that the island is already overcrowded, it could be made into a very wonderful center for biological studies. The possibility that the old volcano might again act up would always be with us, but is it not a dying one, although slowly, if you look back over its history?

Yankee Harbor.

Here there is a great deal of more or less level land upon which one could build. However, the long spit of rock, rounded boulders, stones, and gravel (shingle), strike me as an area that can be terribly wet, wind and wave swept at times. The stones forming the ocean front rampart seem to have been piled up, and rolled about as so many grains of sand in a sand dune. On land there is no protection against storms, although the spit offers some protection for boats within the harbor.

Within the harbor we found the foreshore almost wholly blocked by drift ice, carried there, and held by tidal or other currents. Although a fairly strong wind was blowing across the spit from the ocean side it had no effect on the ice sheltered from that wind by the spit, low as it was. I get the impression that that ice is always here and constantly replenished by slides from the ice cliffs that ring the other half of the harbor.

The better building site, or sites, other things being discounted so far as terrain goes, are on the penguin rookery at the head of the harbor. For this reason, I recommend against this place for a permanent station.

Mosses, a few lichens, and samples of the algae on the beaches were collected. From moss scraped off a rock high on the spit we Berlesed more mites than I had seen so far on this trip. There were some Collembola too in the mosses collected ashore.

#44-63

Two dredge hauls were made at the ship's anchorage in the 30 fathoms. The first brought up a small starfish; a half dozen crinoids, the greatest number of these echinoderms taken in any one place, before this we had but one or two in very few of our previous hauls; there were also a few worm tubes;

several small ascidians; and algal fragments; all were washed clean of bottom material, no mud, no sand, no gravel, no rocks. In the second haul we got a lot of stiff mud from which sponges, many tube worms, ophiurans, ascidians, and algae were obtained.

Our stay was too brief for a trial with the fish trap. These should have not less than an all-night soak; besides, 48 or 36 hours are always better than 24.

One is intrigued with the number of crinoids. More drags could be profitably made, but from a trawler rather than working out of a shore base with a small boat.

I do not favor Yankee Harbor as a laboratory site for reasons stated above -- penguins, wind, weather, and ice.

Hope Bay.

This is an icy, extremely windy place, as Capt. McDonald can tell you.

There are two bases already here, British and Argentine. The latter occupies the most favorable part of the area and has the best installation as well as landing, or dock facilities. Landing from boats at the British Base is difficult except at high tide; a pier once there is no more.

The Argentine Base sits upon what must have been a part of the large Adelie colony nearby. Prospects are that it will, in time, become further depleted as the young pups and sledge dogs (to be) still untrained, run free. While we passed through three young dogs ran down, worried and killed a full grown penguin. How often this happens is anybody's guess. The Argentine who was showing us about went through the motions of pulling the dogs off, slapped at them, which gave us the license to administer a kick or two, all to no avail. The dogs would turn away, but run after the penguin again as soon as our backs were turned. One got in a good bite - there was no use sticking around longer.

With untethered young dogs about one can easily clear off a penguin rookery for a building site. I do not recommend this, nor write this for publication - this paragraph and the one preceding it!

At the Argentine Base the wardroom mess was hospitably entertained at "Asado," a real "criollo" affair.

At the British Base we had earlier had tea. Here we learned that occasionally Chinstraps, and a few Gentoos, turn up; that sheath-bills are resident, skuas and Dominican gulls, well-known; snow petrels are occasional, as are also Fulmers and giant petrels; shags occur in limited numbers; and

antarctic terns (or are they the swallow-tailed ones?) can be found about the "lakes." Young's Point has silver-grey petrels, and Adélies have a rookery on Vorter (sic) Island further south; still farther on one encounters Emperors.

High tide curtailed our dip-netting along shore. Moss was about in very small patches, widely scattered; did not see any worthwhile "stands." What we gathered and Berlesed yielded a very few insects.

The dredge haul at the ship's anchorage, in 25-30 fathoms, mud and sand bottom, turned up many tube worms, some small claims; a gasteropod, ophiurans and algae.

Twenty-two fish were trapped, a goodly catch where there are fish in plenty there must be an abundance of food for them. Stomach contents of those picked out for preservation I expect to have examined in Washington. Having no facilities for curing for a lot of pickled fish, the fish I have saved have been frozen. Let us hope that we can get them all back in the same state.

I cannot recommend Hope Bay as a possible or future station site. I am sure it is located in a biologically rich and interesting area, but as already said, it has been taken over by others. On the score of wind and weather, too, I cannot rate it high in any list of station sites.

Seymour Island - Snow Hill area.

Seymour Island has about as large or perhaps larger snow-free area as any place we have seen to date. It is a soft and muddy place; one could almost call the soil, loam, so soft is it, but nothing much grows on it. Saw and gathered very little plant material ashore.

"Amidships" toward the south is a great "alluvial" river valley with running water and tributary streams, some, though, at this time of year were much reduced in volume.

Few birds, fewer seals, about. One dead one, very old carcass, only skin and bones left, was found in a gully toward the south end of the island.

The island is a barren looking place so far as animal and plant life was concerned when we were there.

Paleontologically, the island has quite a different "face." Much has been done here in this line. I do not know how much more remains to be done. This would bear looking into. Mr. Berg, and the Commodore, in a brief space of time secured a number of highly interesting fossils, wood, bone, and shell specimens.

Most zoologists are interested in paleontological doings; for them paleontology is but fossil zoology.

As to recent organisms: From the ship lying-to in the ice off Seymour Island a few hours on February 15, two hauls were made in 38 fathoms over a rock, sand, and gravel bottom; because they were made at the same place and were so similar, the results were treated as one. Thin, thread-like worm tubes were in this drag by the thousands, along with some clams, and

nemertean worms; included in the haul was our first living barnacle; some alcyonarians; and several stalked, white ascidians that had much the look of "Jack-in-the-pulpits."

Another dredge haul was made while at anchor off Snow Hill Island on 17 February, in 14-15 fathoms, on a bottom of mud so tough and stiff that it was like handling silicon putty, or a tough synthetic rubber mass. In this haul we found a strange polyp, the type of coelenterate represented I do not know; there were also a nemertean worm, three species of annelids; several sea-urchins in part fragmented; some bivalve mollusks; and the first two cumaceans so far seen (cumaceans are small, often tiny shrimp-like crustaceans, bottom feeders).

Two other dredge hauls combined, were made in 12 fathoms over a tenacious mud-cemented sand bottom on the 19th. This time we got more worms; a starfish; hydroids; a piece of an alcyonairian; one tiny amphipod; and a number of small living clams, and some dead shells.

Perhaps the most interesting of all specimens that were secured in this area were three large, red sea-spiders, pycnogonids. These had hopped onto a fish trap let down over the side of the anchored ship on the night of the 18th, supposedly a fathom or two off the bottom so that the trap would not drag as the ship swung with wind and/or tide. The trap hauled up the morning of the 19th held nothing else of consequence, a fragment of a yellow sponge, and scraps of algae.

Tried this "stunt" the night of the 19th but lost the trap as some ice cake or berg coming alongside during the night carried it off. There

seems to be quite a bit of drift ice in desultory movement in this area, probably more wind and current driven than otherwise; but its behavior could not be foretold or counted on.

We really have a rich and promising bottom in this area, in and about Seymour, Snow Hill, and nearby Vega, Cockburn, and other islands. In almost every dredge haul made along the Palmer Peninsula we got some form of animal life that we had not turned up in any of the hauls preceding the one in hand.

It does seem to me that a thoroughgoing marine biological survey be made of the territory - the sea and bay bottoms sailed over in this cruise, in the area where the Eltanin cannot or should not operate.

And for this a vessel or vessels should be provided with all necessary "gear" and equipment, enabling it to work in depths upwards of 400 and 500 fathoms, and with auxilliary small crafts aboard for inshore, shallow water work.

Welchness, Dundee Island.

Here we have acres of snow-free land, enough or almost enough to set up a second McMurdo, but the place apparently has been taken over by the Argentines, judging from buildings, survey stakes, and tractor tracks.

Landing with or from the LCVP was troublesome, beached ice cakes lined the shores of the peninsula or spit that forms the major part of Cape Welchness, leeward as well as windward sides.

Ashore we covered the open, more or less level, snow-free land from one side of the peninsula to the other. Found only scanty and scattered small patches of moss. Apparently few birds about, other than a small flock or group of terns where the peninsula met the ice cliffs on the windward side, none of the birds seen were nesting; Gentoo penguins were few and far between; a Dominican gull or two was flying about, more were "roosting" on an ice cake just off shore. Standing on a small hillock was a skua and her nearly mature chick, but no others in the air at the time. The garbage dumped from the ship always brought more birds around than we saw on, or over, land, excepting, of course, penguins.

Some live seals, not many, were lying on the beaches, more were farther out on ice cakes, but well inland and scattered far and wide over the whole area were 20 or 30 (perhaps more) remains of dead seals -- I would judge adolescent or half-grown, may have been younger. The skin, and the bones within, were about all that was left; heads seemed to have been picked clean. The cause of death - I find it difficult to believe that they could have been stillborn young, much too large.

#53-63

A forenoon and afternoon dredge haul were combined and treated as one; were made in 30 fathoms at the ship's anchorage, mud bottom. As on such bottoms, worm tubes formed the bulk of the catch; a half-dozen species of annelids were represented; one small sipunculid worm; starfishes, ophiurans, and the largest holothurian yet taken; there were also a few coelenterate animals, stalked ones attached to small pieces of rock; and two long rope-like strands of colonial ascidian. Another of these, $1\frac{1}{4}$ -inch in diameter "ropes" was brought to me by a member of the ship's recreation party that was put ashore in the late afternoon. Also a dried pycnogonid (decalopoda antarctica) was picked up on the beach; as well as a specimen of "krill" from near the mouth of a small melt-water stream flooded by the tide. Of the great windrows of algae on the beach a few samples were saved.

Here we lost the two traps that we have been using regularly of late. So many and such large icebergs moved in on shore that we could not locate the trap floats. We still had one trap in reserve and the engineering department on the Staten Island quickly constructed us another out of wire that I had purchased in New Zealand for just such a contingency. But there was no opportunity to undertake a second trial here.

Roughly, the biological potential hereabouts, from the marine biological point of view, rather good or better, bird and plant life are scanty.*

* I think Lenton report remarks that some marine biological work was carried on here in 1951-52. Shall check.

There are fossils to be found. We got a fine specimen of an ancient clam. Unless the paleontology here has been covered by others, there still may be an opportunity to accomplish something in this line at Welchness.

The apparent preoccupation of the site, by Argentina, however, does argue against a U. S. biological station here at this time.

False Bay, Livingston Island.

Weather none too good, had to wait it out before we could be landed ashore even by helicopter; no place in this weather to beach shore boat; surf or swell breaking on the steep-to shingle and boulder covered beach (moraine debris?) rendered landing by boat impossible! - at least in this, and I would say in much other weather either. Getting in and out, off and on shore is a must for any station, and this should be possible most of the time. If not, the place is out as a station site.

#60-63

Did gather a fair collection of lichens and moss; Berlesed the latter aboard. Not time enough spent here to warrant setting fish traps.

Chinstraps were the only penguins seen ashore, a couple of rather widely separated, lonesome little groups of a dozen or so sitting forlornly among the rocks. No evidence that there had been a rookery on this beach. These penguins were probably of this year's crop "on their own," having been cast out at home; some of them were half through moulting their "chick" plumage.

Around the "corner" in South Bay, Commodore Lewis spotted a number of seals resting as usual, but this time in a fresh-water pool. He said also that there was a colony of penguins about 400 feet up, probably Chinstraps, those are the only ones we saw elsewhere in these parts - and how the little "beggars" like to climb high! As his helicopter swooped down for a closer look, the penguins all ran toward the edge of the cliff, several dived over, to be killed on the rocks below. Seeing this, Commodore Lewis did not continue flying in this neighborhood.

#61-63

In the dredge: muddy, coarse sand, rocks and gravel from 18 fathoms along with hydroids; no end of worm tubes of various sorts; two ophiurans; mollusks; an amphipod; two isopods; one specimen of krill; one pycnogonid; some long strands of a colonial ascidian; and a few solitary stalked ones.

Might enjoy dredging here, but a station in this place - NO!

Discovery Bay, Ash Point, Greenwich Island.

Quite a wide open place, windy as all "get-out" while we were about, full of shoals; weather thick, to say the least.

Went out with survey party to Ash Point, hoping to be able to pick up our fish traps that had been set out the night before (February 25), but were recalled to the ship before this could be done.

Lt. Beam, who went ashore with the survey party, undertook to check on the penguins there. He saw no more than five or six Gentoos, and a single Chinstrap.

At this time of year, and in the weather we encountered here, the Point is a bleak, sparsely, if at all, populated place. In general, Chinstraps seem to predominate in the rookeries so far seen in the South Shetland Islands, I expected to find more here. Still, we did not get around as anticipated because of the weather. A number of the ship's personnel got stranded ashore, and an equal number of Chileans spent the night aboard the Staten Island when the seas and wind got too high the evening of the 25th.

#62-63

A dredge haul on the 26th, in 31 fathoms, mud bottom, at the ship's anchorage in the Bay turned up the first sizeable brachiopod of the cruise. Although no larger around than a nickel, it was ever so much larger than the few smaller than pea-size specimens we got on just one other occasion; otherwise, there were a lot of tube dwelling worms; three or four species of hydroids; bryozoans; a dozen extremely thin-shelled snails, mostly broken or crushed; and a half dozen of the rope-like colonial ascidian such as we had taken on several other occasions. This time we got the complete animals,

it seems; I did not know that they had "roots" holding them to the bottom. Well, they have -- good for our old steel drum! I always thought these "fellows" were free-swimming like salps.

#62-63

In the fish traps when we got them back were just two notothoniid fish which I took to be different species, both were saved, a third little orange-colored fish said to have been in the trap by the "boys," got away in the shuffle; otherwise, there was just a starfish, an ophiuran, a small, crushed red sea-urchin; and a few fragments of algae.

Except for the interesting sea life that seems to be had here, the area as a whole does not particularly appeal to me. Earlier in the summer there may be more animal life about, but if we are to biologically do Palmerland, I would rather be closer to it.

Includes
Potters Cove
February 28,
March 3-4, 1963
Mar. 4

Ardley Island area and Potters Cove, King George Island.

This February 28, at 0515, Capt. McDonald and I had a helicopter flight over Ardley Island and vicinity. We did not get over Potters Cove. There were lots of lakes in the hills and lowlands. Had a good view of the reef exposed at low tide that leads some to call the island Ardley Peninsula.

The bad weather we encountered on March 3, that ruled out all operations in Collins Harbor, continued over the 5th and interfered with all plans. The survey party that went ashore, had only 15 minutes there before being recalled.

#64-63

Weather, however, did not interfere with a dredge haul on the afternoon of the 28th. The very muddy bottom over which the drag was made, 47 fathoms down, must be covered, literally, with a forest of worm tubes, the work, I should say, of some 8 to 10 different species of worms; an equal number of ascidian species were counted in this haul too, so intimately must some of them have been associated with the tube worms that they had grown around the tubes, and used them for supports; there were also hydroids; three or four species of mollusks; three of sponges; two of starfish; and two of ophiurans; lastly, a single isopod.

We did get our fish traps over but they were left on the bottom to be picked up on our return from Arthur Harbor where we were to go for the next day or two. It was a generally bad day when we got back to them on March 3.

On our way to get the traps we dropped a survey party, including

Mr. Crowell and Capt. McDonald, on a beach between the Ardley and Fildes peninsulas.

Shortly after we got the traps up we were recalled to the ship, the idea being to give Lt. Beam a chance to look over an old wreck somewhere in the vicinity. He did not get to go nor did we get to take off the shore party. They were left to their fate, so to speak. Notwithstanding, they had themselves a party bonfire and made a show of shelter building in anticipation of a night ashore. Only Mr. Crowell out of the entire group of 8 or 10 had any survival gear with him - no radio, no boat.

At low tide the party probably could have made it over on the connecting reef to Ardley Island where there is a shelter hut, in case of necessity. Even so, they found a food cache right where they were stranded. It had been left behind by an earlier English survey party. However, no need arose to open it. For about suppertime they were rescued.

The water inshore, although we could not see how it was from the ship at anchor, was relatively smooth and quiet compared with the sea out where the ship was "bouncing" up and down. All's well that ends well.

So impressed was Capt. McDonald, who was ashore with the survey party, with the excellence of the area as a building for shelter for boats, and the quiet water inshore, that I believe this Fildes-Ardley Peninsula site became his second choice site for locating the Palmer Peninsula biological station.

As this was no day for the birds and was so very late in the season, little can be said of the bird life of the vicinity. There were seals about. The question was raised, could they be fur seals? There was no chance to investigate.

In the lifted traps, six fish were found, $10\frac{1}{2}$ to $13\frac{1}{2}$ inches in length, half a dozen starfish, and a number of red amphipods.

March 4 - Potters Cove.

73-63 This afternoon ashore, mosses, lichens, clumps of grass, and samples of marine algae that had been thrown up on the beach were collected. Berlesed the mosses; later, when I took moss and grass out of the bag in which they had been placed, a few insects (flies?) dropped in the pan over which I was working, and were quickly bottled.

There were several crab-eaters hauled out on the beach. Gentoo and Chinstraps were to be seen but in very limited numbers, a few of each at most, and fledglings at that. Also noted were Dominican gulls, skuas, Wilson's storm petrels, and two birds that Capt. McDonald said were antarctic petrels, or perhaps giant petrels. Lenton in his report listed sheath-bills, but we saw none.

74-63 The dredge haul of this day, in Potters Cove, in 19 fathoms, mud bottom, produced a fine lot of mud-dwelling tube-building worms, if anything larger and handsomer than those we got off Ardley Island but of the same species surely; the mollusks, likewise, were of the same species as yesterday, but the sponges, hydroids and echinoderms present in the haul of the day before we failed to find; did get an alcyonarian, though.

After a morning flight over the area, Capt. McDonald remarked that Potters Cove seems to have about everything: the necessary piedmont insuring safe landing for aircraft, with space on the snow-free land, too, for an airstrip, certainly for helicopter pads; a dock, mooring facilities, and a

sizeable lake for water supply. The dredging holds forth hope for more rewarding collections than we made during our brief stay here, but I, for one, would like to leave the South Shetland biology and geology to the British who have already done so much in these fields in this general area, and in and about these islands.

For March collecting
see sections II
and III

March 4, 5, 1963

Admiralty Bay.

Here, too, we struck bad weather, high winds and poor visibility, and were unable to effect a boat landing on either side of the Keller Peninsula. Tried the east side the afternoon of the 4th but found too much ice in the shallow water inshore. The survey party was put ashore by helicopter.

Because of continuing unfavorable winds the projected landing on the west side of the Peninsula on the 5th was also given up.

#75-63

However, a dredge haul was accomplished on the 4th. It was a rather meager one, containing a number of tube-building worms, starfish, and a clam or two. The depth was 22 fathoms, the bottom mud.

Our fish traps put over this same evening, but not lifted until mid-morning of the 5th (having waited for the wind to die down), contained a very scanty catch, a couple of worms; some aphipods; and a starfish.

Coming up this area under very unfavorable circumstances, our trials were limited, nor could a thorough examination be made of the site on the each side of the peninsula. However, there appeared to be land enough for an extensive installation if one had to be placed here.

The last we saw of King George Island was Penguin Island off to the east. Between poor visibility and the need for keeping a safe distance offshore in bad weather we were not in a position to identify the bird colonies said to be on this little island and the adjacent nearby shore.

As we passed the spray over the ship almost obscured our sight of the island, sheets of it went right up against the wheelhouse windows, a good 70 feet above sea level.

Thus it was as we bowed out of the Palmer Peninsula area and the South Shetlands.

APPENDIX B

ITINERARY, PALMERLAND SURVEY
(in and about Palmerland)

1963

Jan. 18-19	Marguerite Bay
Jan. 20	Rothera Point
Jan. 21	Under way for Anvers Island
Jan. 22-25	Arthur Harbor
Jan. 25-26	Port Lockroy
Jan. 27	Dorian Bay - Pleneau Island
Jan. 28	Petermann Island
Jan. 29	Galindez Island
Jan. 31	Cape Tuxen - Green Island
Feb. 1	Paradise Harbor
Feb. 2	Paradise Harbor
Feb. 3	Chilean Station
Feb. 4	Danco Island - Couverville Island
Feb. 5	Cape Spigot
Feb. 6	Gamma Island, Melchior Islands
Feb. 7	Gamma Island, Port Lockroy
Feb. 8	Dorian Bay
Feb. 9	Svend Foyn Harbor
Feb. 10	Lay-to in Salvensen Cove - Brialmont Cove, Alcock Island
Feb. 11	Alcock Island
Feb. 12	Whalers Bay (or Bight), Deception Island
Feb. 13	Left for Yankee Harbor
	Left for Hope Bay

Appendix B-2

Feb. 14	Hope Bay
Feb. 15	Left for Robertson Island
Feb. 16	Off Argentine Ice Shelf Base
Feb. 17	Off Snow Hill Island
Feb. 18	North end Seymour Island anchored between Vega, Cockburn and Seymour
Feb. 19	Seymour Island
Feb. 20	Welchness
Feb. 21	Suspiros Bay
Feb. 22	Underway for Alcock Island
Feb. 23	Spring Point, and Alcock Island
Feb. 24	Landed on Spring Point
Feb. 25	False Bay, Livingston Island
Feb. 26	Discovery Bay, Ash Point
Feb. 27	Ardley Island
Feb. 28	Ardley Island
Mar. 1	Port Lockroy
Mar. 2	Paradise Harbor
Mar. 3	Ardley "Island" Cove, King George Island
Mar. 4	Potters Cove, King George Island
Mar. 5	Admiralty Bay, passed Lion's Rump and Penguin Island
Mar. 6	Underway for Valparaiso!!!

APPENDIX A

PALMERLAND SURVEY, 1962-63

Tow-net hauls made:

T-1-63	Marguerite Bay	January 18
2	Bonaparte Inlet	24
3	Arthur Harbor	25
4	Port Lockroy	26
5	Petermann Island	28
6	Galindez Island	29
7	Galindez Island	31
8	Paradise Harbor	February 1
9	Bryde Island	2
10	Paradise Harbor	4
11	Danco Island	5
12	Melchior	6
13	Svend Foyn Harbor	9
14	Whalers Bay	12
15	Whalers Bay	13
16	Hope Bay	14
17	Welchness	20
18	Alcock Island	23
19	Discovery Bay	26
T-20-63	False Bay	25*

* Was numbered wrong, date is correct; will do as is.

Introduction:

Says
center

The following is a resumé of the biological investigations carried out at the sites visited by your Survey party in and about the Palmer Peninsula, Antarctica, and the South Shetland Islands. These are set forth in 18 sections I to XVIII, more or less chronologically and geographically arranged, as follows:

under the heading "Site Investigations,"

- I. January 18-20; Marguerite Bay area—
Adelaide Island, Base T, Avian Island,
Rothera Point
- II. January 22-25; and March 1; Arthur Harbor area—
British Base IV,
Branscombe Point and vicinity, Janus
Island, Forgeson Island.
- III. January 26-27, and March 1; Port Lockroy-Porian Bay area, Base A.
- IV. January 27-31; Argentine Islands area
— Lemaire Channel, Danneborg Islands
Berthelot Islets (British Base is on
Galindez)

- V February 1-3, and March 2; Paradise Harbor
area — including Chilean, Gabriel
Gonzales Videla Base and the Argentinian,
Almirante Brown Station.
- VI February 4, 5; Danco - Couverville Island
area
- VII { Melchior Islands area;
February 6, 7;
- VIII February 8-10; Wilhemina Bay -
Svend Foyn Harbor - Solvensen Cove
area.
- IX February 10, 11, 23 and 24; Brialmont
Cove, Alcock Island, Spring Point.
- X February 12, 13; Deception Island
- XI February 13; Yankee Harbor
- XII February 13; Hope Bay.
- XIII February 15, 17-19; Seymour Island -
Snow Hill area.
- XIV February 20, 21; Welchness, Dundee
Island.

XV February 25; False Bay, Livingston Island.

XVI February 25, 26; Discovery Bay - Ash Point, Greenwich Island.
(For Yankee Harbor, this Island, see section XI February 13.)

XVII February 28, and March 3, 4; Ardley Island area, and Potters Cove, King George Island.

XVIII March 4, 5; Admiralty Bay.

¶ All listings are of a very general nature. It was not possible to enumerate or identify all the seals and birds in any one area or the animals taken in the dredge or fish traps, the insects or the species of cryptogamic plant life.

The dredged material by force of circumstances, want of time and neces-

sary equipment for handling it, was not worked over as thoroughly as I would have liked. What has been saved in the way of specimens collected by dredge, tow net, and otherwise must needs be identified by specialists conversant with the various groups of organisms before their geographic distribution, their relation to their environment and their associates can be discussed, and before our findings can be compared with those published by previous expeditions which worked over or collected in the same areas. A list of tow net hauls made is appended to this report.

~~Nevertheless~~ ~~However~~ as annotated, it is believed that these listings will convey some idea of the nature of the flora and fauna, terrestrial and aquatic, at, or in the vicinity of the sites investigated.

⑤

The comments, recommendations, and conclusions which follow are my own based on personal observation, and on discussions with other members of your ~~Survey~~ ^{Party} Party.

Caps
center

In Conclusion

Though terrestrial animal and plant life was more abundant in some areas than in others, no truly barren area was found in the sea.

Seals and their numbers are mentioned when they came to our attention. Some effort was made to note bird rookeries seen by various members of our party and by ship's personnel, and where possible, which was not as often as we would have wished, to identify the species inhabiting them.

Noteworthy bird colonies, mostly penguins have been noted in the ^{area} sections of his report ~~in the areas covered by him~~ (all but sections

^{XIII} VII, XIV, XVII, XVIII, in which poor visibility, insufficient time for searching or lateness of season ~~might~~ may have kept us from discovering any rookeries of consequence.)

Each and every area visited in the course of this Survey holds great promise for ~~the~~ ^{marine} biologist working on, or with particular ~~marine~~ animals or groups of them. Representatives of most, if not all phyla of marine animals, even though we may not have turned them up in our samplings, are sure to be found in the regions explored by us.

Ice conditions more than anything else will restrict, at times, the operations of those who may follow us, as such conditions hindered us on several occasions. But if we wish to know how Antarctic and Subantarctic organisms live, and where, and under what conditions, and above all know their kinds and distribution we must not fear ice, or seek wholly to avoid it. Indeed, to get a true picture, and the whole story concerning any and all of them, we need to go where ice is

Though it may add to the difficulties of planned or to be planned operations.

The word "hazards" might have been used along with difficulties but hazards will always be minimal if one goes with proper and adequate equipment and possessed with, or accompanied by some one with "know-how" born of experience. Do not be too fearful of

"ice" in choosing the site ^{of} ~~for~~ your station ^{or your studies}
~~over~~ ^{over} ~~the ice~~ ^{here} ~~here~~

Personally I favor locating the station as close to the mainland of Palmer land as possible and practicable [This was to be a Palmerland station was it not?] and not in the South Shetlands. Leave the work to be done in those islands to the British who have already done so much, and published so much on that area.

I also favor a sites or ^{areas} ~~sides~~ from which can be reached with greatest

enter
make as a foot note on page

⑩ Don Squires of the U.S. National Museum who is engaged in working up the corals of "The Ross Sea Fauna" for the New Zealand Oceanographic Institute, is strongly of the same opinion. So far as it is humanly possible we should "pursue" our Antarctic organisms through the winter as well as during the few summer months. The occasional open water that occurs in Arthur Harbor and perhaps more open in Paradise Harbor in the winter months could well render such a pursuit rewarding

turn back
to p. 7.

⑧
facility, and ~~encountered~~ ^{on foot or by boat,} within reason-
able distance, the widest range of types
of environment, and terrain, and the
greatest variety and numbers of forms
of life. ^{animal and plant life}

^{with the foregoing in mind}
I ^{definitely} favor placing the biological station
or stations, if there is to be more than one
or a main base with one or more sate-
lite bases, one or all, on a line running
from Arthur Harbor through Port Lock-
roy to Paradise Harbor.

Arthur Harbor as a site for a
base or major station comes first
to mind as the best possible site for
on most counts. It is a very favored
area, weather, ice, boat shelter, water
supply, and perhaps also "piedmont" for
air craft landings, abundance, though
not necessarily variety of terrestrial
life (birds' primarily), marine life, and
ready access to a great variety of environ-
ments, ^{within not unreasonable distance} This ready access to a great
variety of environments ~~is not to be~~

holds good for the other two sites on our ~~the~~ projected line of stations, Port Lockroy and Paradise Harbor.

Twenty miles to the South of this "base line" is the Lemaire Channel.

no para

Here we spent the night of January 27 drifting about in this scenically breath-taking area where the mountains are as impressive and at the same time as fantastic or as beautiful in shape and form ~~and indeed as big~~ as any iceberg of imposing size and beauty so far encountered on this journey along the Antarctic coast from McMurdo to Marguerite Bay and Adelaide Island to Arthur Harbor and now to this out standing natural beauty spot.

Enhancing many fold the sheer beauty of our surroundings were the crystal clear reflections of the ~~the~~ in the placid waters of the Channel ~~200 feet deep~~ ^{800 fathoms} majestic mountains rising from the water's

This is a punk sentence shall try to fix in double spaced draft.

edge to lofty,

(10)

A snow-topped peaks, ~~and~~ doffing
from time to time their caps of clouds
thus granting ~~at last~~ the "by-stander" ^{Ter} an opportunity to photograph them full stature. ~~⊗~~

⊗ "in all their imposing grandeur" I wanted to add but dared not. I beg ~~for~~ to be forgiven for running in this "rare" but you would have been moved to feel it not say much the same had you been with us that lovely night of January 27, 1963

Thirty to ³⁵ ~~thirty-five~~ miles to the South you come to Cape Tuxen ^{and} Green Island in the Berthelots, with its ~~masses of unbroken vegetation~~ ("... luxuriant growth of moss nearly 41 acres in extent by far the largest unbroken [sic] patch of vegetation yet found in Antarctica". On the way having passed Pleneau and Petermann Island, ~~and the Argentine Islands~~ with their ~~Adelie and Gentoo rookeries~~ ^{shag and tern colonies} and the Argentine Islands, where on Gulindez the British Base is ^{would} have ~~passed~~ ^{seen} through fields of ice on which

to their snow^{-topped} ~~capped~~ peaks wreathed
in ~~drifts~~ '.


numerous seals were lying about. (11)

8/ Eight miles from the Western end of the base-line is Cape Monaco (not seen) with its ~~reported~~ ^{reported as} garland of six islets ~~just off~~ crowded with penguins in season, and 42 miles out of Bismarck Strait you come to Victor Hugo Island, and beyond that the open sea.

9/ Some 50 miles to the North via the Neumayer Channel, an interesting stretch itself, are the Melchior Islands off the North Coast of Anvers Island with its ~~several~~ in hospitable embayments. In the Melchior Islands on Lambda Island is an ~~Arg~~ unoccupied Argentine Base. Though there ^{was} not much on land to interest us, the dredging ~~showed~~ promising. (The fishing was good and)

~~To the Northwest~~

11 Northeastward



~~On the way to the~~ by way of the Errera Channel one passes Danco Island and Couvreville, and 45 miles from Port Lockroy, Cape Spigot with perhaps the second best (^{second} largest) Chinstrap rookery (next to Alcock) in all Antarctica and 5 miles beyond Cape Anna with its reported bird colonies, not seen and not identified as to inhabitants because of poor visibility.

Within 50 miles of the base line, or within a like radius from each of the recommended sites one can have everything in the way of environment animal and plant life that can be had in Palmer land, ^{all} that any biologist might crave for study ~~in~~ in or out of the laboratory. (terrestrial, or marine, or)

One hundred and 10 miles away to the Northeastward also is Brialmont Cove and Alcock Island upon which is perhaps the greatest penguin colony in all Antarctica — all Chinstraps which I hope will go unmolested for all time to come. He who wants

to study Chinstraps can do so on Deception Island where there are several hundred thousand, on Cape Spigot (Nunatak Negro) where even more have their home, or in ~~the~~ Paradise area.

Paradise Harbor I rate as desirable for a land based station next to Arthur Harbor and Port Lockroy third. Paradise Harbor has lots to recommend it, a good site though at present occupied by an inactive Argentine Station - "Almirante Brown, proximity to rookeries of three different species of penguins (so reported) though we only came upon two of them in our few days in this area, good fishing and a promise of richer dredging returns ~~than we had from~~ ^{than we had from} the two samplings we had. Here perhaps more than elsewhere it may be possible to study or

(14)

at least investigate the life and behaviour, distribution, condition and survival of the marine inhabitants of the Harbor, pelagic, littoral and benthal as the tidal and other currents often, if not through the winter maintain enough open water channels to permit open water channels. ~~An~~ attempt ~~is~~ in a suitable vessel should be made to prove or disprove this contention

- 1 Port Lockroy deserved better than a third place rating. It hurts to put so lovely a place so low, especially in view of its well free harbor in which a vessel might safely be wintered (I shall have to ask Capt. Mac about this), and where we found the littoral faunal so very rich. But Paradise Harbor I feel quite strongly has the far greater biological potential.

Within Arthur Harbor, I favor the (15)
British Base ~~that~~ area or site over Bona-
parte Point, or the so-called Bravo
site for reasons stated elsewhere in
this report, particularly in the discus-
sion of the Arthur Harbor ~~Area~~
(Section II, ~~pages~~ pages II 4-II 6.)

I would urge that we endeavor to
get full and clear "title" to all three
places by transfer, as gifts, or by some
reimbursement, permitting the present
builders/^{or owners} of the huts in situ in Port
Lockroy and ~~the~~ Paradise Harbor to
remove buildings and contents should
they care to do so. We should by all
means start anew, and with our own
new buildings, or structures.

And now? What are we wait-
ing for? Let's go!

start
SEP sheet

Heading here → Appendix B

(centered)

Itinerary, Palmerland Survey

(in and about Palmerland)

1963

Jan. 18-19 Marguerite Bay

Jan. 20 Rothera Point

Jan 21 underway
for Anvers Island.

Jan. 22-25 Arthur Harbor

Jan. 25-26 Port Lockroy

Jan. 27 Dorian Bay - Plonear Island

Jan. 28 Petermann Island

Jan. 29 Galindez Island

Jan. 31 Cape Tuxen - Green Island

Feb. 1 Paradise Harbor

Feb. 2

"
Chilean Sta. ""

Feb. 3

Panco Island - Convergence Id.

Feb. 4

Cape Spigot

Feb. 5

Gumma Island, Melchior Ids

Feb. 6

Feb. 7

" " , Port Lockroy

Feb. 8

Dorian Bay

Feb. 9

Svend Foyn Abr.

(Lay to in Hughes Bay)

Itinerary

Feb. 10	Lay to in Sulvenson Cove - Brialmont Cove, Alcock Island.
Feb. 11	Alcock Island Left for Deception Island
Feb. 12	Whalers Bay (or Bight) " "
Feb. 13	Left for Yankee Abr. Left for Aoape Bay
Feb. 14	Aoape Bay
Feb. 15	Left for Robertson Island
Feb. 16	Off Argentine Ice Shelf Base
Feb. 17	off Snow Hill Id.
Feb. 18	N. end Seymour Island anch. between Vega, Cockburn and Seymour.
Feb. 19	Seymour Island
Feb. 20	Welchness
Feb. 21	Suspiros Bay
Feb. 22	Underway for Alcock Island
Feb. 23	Spring Point, and Alcock Id.
Feb. 24	landed on Spring Point

Feb. 25 False Bay Livingston Id.
Feb 26 Discovery Bay, Ash Point
Feb. 27 Ardley Island
Feb. 28
Mar. 1 Port Lockroy
Mar. 2 Paradise Harbor
Mar. 3 Ardley "Id." Cove. King George Id.
Mar. 4 Potters Cove, King George Id.
Mar. 5 Admiralty Bay, passed Lins
Rung and Penguin Island
Mar. 6 Underway for Vapo! !!!

center cups

~~Postscript~~

regarding

Here on

center
cups
over

{ A few questions and comments ~~here on~~
regarding biological work in the
Palmer Peninsula ³¹ by ~~holder~~ L. Schmitt ^{P.D.}

Looking back over our site survey, what
specifically is it that USARP hopes to accomplish
with the proposed Palmerland laboratory, or
"boat", either one, or both?

Are these facilities to be open, available, and
unrestricted for each and every applicant for
desiring to ~~the purpose of~~ engaging ~~any~~ in some phase, major
or minor, of any line of research that he
might fancy in the broad field of biology?

Or does USARP, to begin with, wish first
to further investigations directed toward care-
fully considered, definitely stated objectives
— researches holding forth promise, or at
least the hope of yielding information being
sought, or of solving specific biological
problems concerned with Palmerland's in-
habitants, plant or animal?

Res. Department in Italy
Enclaves
US & Mexico

In fact, in order to determine what those facilities should consist of, does not something definitive in the way of a program or plan for the Palmerland operation need to be formulated, and made known in advance, at least for the first year or two of their existence?

Otherwise, could not a lot of effort be expended in all, and perhaps wholly unrelated directions in diverse investigations that might never contribute toward the building up of the substantial, well founded, interrelated, cohesive body of biological knowledge which should result from the research work that the National Science Foundation may be called upon to underwrite in Palmerland?

Should not "first" things come first?
 "The first need," as set forth for the Antarctic as a whole in the Projected

Antarctic Science Program by the Chief Scientist of the Office of Antarctic Programs, Dr. A.P. Crary, "is in the field of taxonomy and systematics, the classification and description of the flora and fauna, not only of Antarctica but also of the many islands of the high southern latitudes and even the ~~southern~~ southern mainlands of other continents. Along with his description should go ecologic studies of the environmental and climatic factors under which the flora and fauna live...."

Do we agree that these things should come first in any program set up for Palmerland biological studies?

If so, is not a comprehensive, ~~promising~~ thorough-going biological survey of an area representative of a variety of environments and a diversi-

ty of species, in order?

Such a survey would reveal the distribution of types of bottom, the kinds of animals and plants living on them, the communities and associations they form, their abundance and seasonal occurrence, as well as the "flow" and the constitution of the waters which distribute their larval stages, and in which still other animals and plants drift about, if they are not free swimming forms.

How best to attain these ends — a shore station or stations out of which to operate with a small craft, thirty feet in length or less, as was attempted at the ~~Wilkes~~ ~~Bullet~~ station, I believed[?] for by a sea-going trawler type of vessel, able enough to travel back and forth from a suit-

New para
begin with
Cup. O

~~Some~~ port on the South American mainland, or even the States, if need be, equipped to explore depths of from 450 to 500 fathoms, capable of working in, or at least about icefields, and provided with laboratory space and accommodations for a scientific staff of four, perhaps.

Such a vessel could also work the intermediate and surface waters ~~by~~ by means of its auxiliaries shallow inshore waters, and if, and when necessary support "excursions" ashore for related studies, fresh water and terrestrial. ~~the~~
~~But~~ This vessel may need a shore base, or supply depot conveniently located ^{or near} in the area to be surveyed.

In no way is it intended here, to minimize or overlook what can be

(6)

done land-wise from a shore-based laboratory. But could not microbiological studies of soil, air, and of water and studies of bird and terrestrial plant life be conducted as well and to the same extent from a vessel as from a shore station, if the be deemed of such vital importance that they must be initiated the very first year or two ~~at~~ ^{is} this vessel ~~is~~ operated?

Penguin and other ornithological researchers ^{otherwise} have ~~perhaps~~ the opportunity of pursuing such studies, by living, as ~~the~~ observers in one ~~at~~ or another of the presently active bases in ~~Antarctica~~ Palmerland, Argentine, Chilean, and English. The little studied Chinstrap penguins can be studied at close range, as well ~~at~~ Deception Island, and in Paradise Harbor as ~~well~~ they can be in (or at) any of

the colonies of these birds observed in the course of the survey cruise.

The prospect of seeing, hearing, or reading about what may come out of a biological survey of the channels, straits, bays, and harbors of Palmerland is truly a fascinating one.

N.B. "The Fauna of the Ross Sea" published by the New Zealand Oceanographic Institute is based on the biological materials gathered by that Institute's oceananographic survey of the Mc Murdo area, so to speak. It includes "records" established also by earlier British and Commonwealth expeditions, and occurrences of species previously collected in Palmerland by these and other English expeditions.

U. S. A. R. P.

center Biological Investigations

center Palmer Peninsula and

center South Shetland Islands

center

1962 - 1963

Little Gage

I have separately
paged sections so that
they can be handled individually
as sections and this can be
added to if (necessary) without
upsetting any pagination.

Introduction however can
be page numbered 1 to —

5 carbons please

Behm

Note:
On a separate sheet other
wise blank put:

Caps
& center

Site Investigations

~~UD~~

~~PRX~~

XV February 25; False Bay, Livingston
Island.

XVI Discover-

Start each
sect
sep. sheet

(I) January 18-20, 1963 ^{I ①}

Marguerite Bay area — Adelaide Island,
Base T, Avian Island, Rothera Point.

The ice fields through which we passed getting here had quite a number of seals on ice cakes here and there. Mostly if not all Weddells I take it. Some said they saw a leopard seal but I saw none that quite fitted the description given.

Seals must be plentiful in this area; it was said that about 500 a year are killed for dog food and station mess.

At all places the "usual" birds, many still nesting here: Adelie penguins, Skuas, Black-backed (Dominican) gulls, Blue-eyed shags, and a giant petrel; At Rothera Point also, Antarctic terns.

Mosses and lichens were collected at each place; from them mites and spring-tails were Berlesed; got about three different species of insects.

From a fresh-water pool on Avian Island a few fairy-shrimp (*Branchinecta*) were dipped with an empty beer can lying about (Tom Berg coll.)

Fresh water and marine algae were collected; and at Rothera Point red and green snow algae as well.

From fish-trap set in 50ms off Avian Island, no fish, a number of nemertean worms amphipods, 2 species of starfish; 3 small sea-urchins these last at Rothera Point where the worms, and amphipods also appeared in the traps.

Dead limpet shells left by the gulls whose principal food they ~~limpets~~ seemed to be were scattered about everywhere some with serpulid worm shells attached.

There was much ice floating about making tow netting difficult. This was said to be the best year in the last six, more open water than usual. However the ship had to shift position to avoid a sizeable ice berg; can be a nuisance here.

Avian Island despite its teeming bird life, fresh-water pools, many mosses, is not to be recommended as a station site because of the interference with the penguin rookery (and the "mess" it would have to be

located in; no suitable space not occupied.

Base T site hilly; good buildings, best is the one man "weather shack"; ideal set up of what a one man study and laboratory could be; sleeps in.

Rothera Point, though has the building land - acreage and "Foundation"-wise, a fine place; a low level saddle between the "hills" toward the Point and the more mountainous piedmont to the right as viewed from the Bay landing. I wonder how wind swept this place is; looks like a "draw". We had a fine day here, only a windy and stormy one would tell.

There were only a few penguins here compared with the hundreds at Avian island. Lt. Thomas attempted to walk around the shore of the Point but gave it up because of the ^{determined} way the Skuas which must still have been nesting, dived at him; thought he saw evidence of what might have been a penguin rookery but no birds on it. Quite a few small bergs about in the landing area; seals on a number of them.

(II)

January 22-25
March 1, 1963

Arthur Harbor area → British Base N,
Benaparte Point and vicinity, Janus
and Dorgeson Islands.

There were quite a number of seals about; we had no difficulty in locating a Weddell on an ice cake for bait.

The area is rich in bird life, especially Adelies. There is a large and populous rookery on Dorgeson Island, and other sizeable colonies scattered through the area. About Cape Monaco to the West are 6 islets with rookeries on them. There is also a colony at Biscoe Bay on the S.W. coast of Anvers Island (fide sailing directions); others on Litchfield, Halfway and Humble Islands (fide Berg helo flight), Giant petrels, Skuas, Dominican gulls, as always and in considerable numbers, Antarctic Terns, and Sheath-bills. Many dead limpet shells ashore, others gathered from rocks at tide level.

Mosses and lichens, grass clumps collected, and insects Berlesed from this vegetation.

Fresh water algae and fairy shrimp were found in the pools atop the bluff above the

Base N. hut. Along shore at Bonaparte Pt. and at Base N. marine algae and a few amphipods were dip-netted.

#7-63 Dredge hauls were especially productive - a varied and abundant fauna on a stiff blue mud bottom, 12-17 fms. - a species of sponge, sea urchins, 2 sp. starfish, worms in number, the nemerteans proved to be the common species throughout the area (Amphiporus), 2-3 species of annelids (Nereis, Terebella and at least one other), 4 species of mollusks, and several species of amphipods, an isopod, and 4 or 5 species of sea-squirts (ascidians)

#67-63 The dredge haul of March 1, on our return to Arthur Harbor was a repeat of the earlier one with a host of annelid worms and mollusks.

#8-63 The fish trapping was equally rewarding 88 notothoniid fish were taken in two sets off Janus Island together with starfish and nemertean worms. The fish ranged from 7 to 14 1/2 inches in length. One trap set in the rocky part of Bonaparte

Inlet was crushed under a small berg that either drifted in with the tide or possibly was calved off during the night from the ice cliffs lining one side of the head of the inlet; considerable sea weed and a large clump of ascidians were dislodged as we got the jammed trap up after some ^{hours} labor, tried in forenoon, were successful with a grapnel in the afternoon.

Brash-ice troubled our March 1 landing at the Base N site, recent snows had whitened the ground and the large lake a few hundred yards behind the hut was thin-ice covered, as were the freshwater pools on the top of the bluff above the hut. A few lonesome looking Adelies were standing on the sea-ward side of the bluff; Skuas were about but in lesser numbers than in January. Where do they go; migrate as do the penguins? Porgeson Island was all but deserted, just a scattering of penguins about.

The approaches to Arthur Harbor because of the current running through it, is said to be quite free of sea-ice more or less the year

round.

Regretably during the January two hour squally we experienced in Arthur Harbor we did not get to see the Inlet; it is a remarkably protected quiet bit of water. On March first like a mill-pond, though it was windy and rough out in the harbor.

I do not particularly fancy the Bonaparte Point site because of its exceeding rockiness - large jagged, broken rocks, and a rather uneven gullied (or is it ridged) terrain. The Inlet though a piece of very quiet water beautifully protected from wind and wave might be a very dangerous place. It narrows down to about 80 feet in width, and on the right hand side are ice-cliffs of about that height. These have plenty of fissures, cracks, and crevasses. A fall of a mass of ice and snow could well spell disaster to any vessel that might find itself in the Inlet at the "right" (or would it be the wrong) time. Really, how very accessible is the piedmont here?

Getting fresh water would be more of a problem than at Base N where it could be conveniently piped from the the large and quite deep fresh water lake there. No dam would be needed the valley in which it lies could hold many times the present amount of water without damming; or much of one; but the supply of water seems ample for the needs of any station that might be ~~placed~~ established here.

within walking distance and not ~~too~~ ^{enough} near ~~to~~ be objectionable, ~~to~~ one side is a penguin rookery, on the other the bluff on top of which are situated the fresh water pools already referred to. A man needs to get about a bit for relaxation if for no other reason. At Bonaparte Point he could walk about on rocks and still not "be" or get anywhere.

True the boat landing is not of the best but it could as easily be im-

proved as that, that would have to be provided at the Bonaparte Point site.

During the January squall the Navy survey party was caught out at the Base N. landing. Lt. Nash said it was "not so bad" and that there was some degree of protection from what went on in the harbor outside.

I might add here too that Lt. Nash also favors the Base N. hut site over Bonaparte Point from an engineering point of view. I have his permission to so state.

On the 21 January we were under way for Anvers ~~Island~~ Island and Arthur Harbor.

III

January 26-27
March 1, 1963

Port Lockroy, Dorian Bay area, Base A;

In this area the Gentoo penguins take the place of the Adelies but for this major difference bird life is as varied as elsewhere though the number of individuals seems less than at Arthur Harbor. I do not recall seeing any Giant petrels here. Rookery space is more limited resulting in fewer colonies, but those available are well populated. Near at hand to, but not on the Base A site are sizeable Gentoo colonies, and also at Damoy Point on the way to Dorian Bay is another large one. A colony of terns holds forth on a rocky, steep headland near the entrance to this Port, and a well populated shag colony occupies an island in Peltier Channel (Priest Id?) on the way to South Bay, Doumer Id and where the Chilean Base "Yelcho" is located. Additional colonies of Gendbos occur down this way, also. Skuas, Wilson's storm petrels and sheath-bills were about too, and in the Harbor.

Three
chins straps
were seen
among the
Gendbos
here by
Lt. Thomas.

There was comparatively little vegetation to be seen, nothing at all to compare with the luxuriant vegetation that we were to see later in the Argentine - Berthelot Island area. From moss gathered here a few Collembola were Berlesed.

#9-63
#66-63 The more meager dredge hauls here, over a muddy bottom, in general character resemble those made at Arthur Harbor. Worm tubes were abundant; ophiurans (serpent or brittle stars) and ascidians of species other than those obtained Port Arthur were washed from the mud. Additional specimens of these species were secured by the same means on occasion of our March 1 visit; and unexpectedly a tiny gastropod.

#10-63 The richness of the shore fauna is worthy of special mention. It so happened that we landed at the Base pier at extreme low tide. In the course of a few scrapings along the rocky shore Poblete's dip-net brought up a variety

animals — the already (to us) common nemerteans (Amphiporus); three sea-anemones resembling the North Temperate Metridium (the only sea-anemones of the trip it turned out), two species of amphipods, and two widely different and very characteristic Antarctic isopods, Serolis and the giant, almost 10 cm. long Glyptonotus antarcticus. Had we had a favorable tide in the Arthur Harbor area, I like to think that we would have been as fortunate there.

#11-63 From the fish traps we got more nemertean worms than we knew what to do with, 2 species again of amphipods, and 7 nototheniid fishes.

The building area about Base A (Goudier Island) is a rocky one but no where as rough going as on Bonaparte Point, Arthur Harbor. Port Lockroy is one of the few swell-free Palmer land harbors. As to air-craft facilities, landings, see Capt. McDonald.

I cannot refrain from adding that Port Lockroy is as scenically beautiful a place as I could spend an Antarctic summer in, or a winter too for that matter — Lemaire Channel is another.

IV

IX(1)

January 27-31,
1963

Argentine Islands area — Lemaire
Channel, Danneborg Islands, Berthelot
Islets. (British Base is on Galindez)

Lemaire Channel, Jan. 27-28, Pleneau Island,
Jan. 28, Petermann Island, Jan. 28,

Galindez Island, Jan 29, Cape Tuxen
and Green Island, northern most of
the Berthelots, Jan. 31.

Galindez
Base
gets its S.W.
from a large
pond (lake?)
behind the station

Many seals were basking on the ice cakes,
thick in all directions, south of Galindez,
more seals over a larger area than any
in which we had seen any large number
of them. Two whales were seen north
of Galindez on the 29th

An interesting and well populated area,
animal, as well as "vegetable"-wise. Many
birds breed in this area. The following
list was given to me by Mr. R. Lewis of
the British Base here on Galindez:

Dominican (black-backed gull) — a year
round resident, an interesting record; Great
Skua ^(breeds) in Antarctic, or is it the Swallow-Tailed
Tern? (breeds); Giant petrel (visitor only)

always seems to appear in bad weather — several to half-a-dozen of them), Shear-bill (winter visitor); Wilson's storm petrel (breeds); Snow petrel (breeds locally) Adelie penguins (are on Dalour and Petermann Islands), Gentoos also on the last named island; Chin straps (casual visitors to the area), Emperor penguin (seen once); Blue-eyed cormorant (shag) (breeds). A cormorant colony not seen by us is located on Winter Island.

On Petermann Island both Gentoos and Adelies have rookeries; among the Gentoos one Adelie made itself at home. On a rocky peak across a snow field behind the Gento rookery, Antarctic terns were flying about in considerable numbers, some were nesting. Jack Crowell saw one with a chick (close up — beauty of this nesting site was the fact that you could walk on to and around it to photograph the birds close up.)

At Cape Tuxen there is a colony of

41 blue-eyed cormorants. On our return to the ship from Green Island we tried to get photographs of a flock of 30-40 (? 20-30) of these shags sitting on the water quietly and in a surprisingly close group, but they proved ^{before} wary; ~~as~~ we got into comfortable camera range, they dived and scattered. Green Island has a considerable population of Skuas. Judging from the aggressiveness displayed by several of them, they were protecting nests and young. A large colony of cormorants is also located 4 miles S. of Cape Tuxen in the Berlesed Ids.

Mosses and lichens were gathered from the several islands and the Cape, and Berlesed. An amazing ^{number} of Collembola were driven out of a couple of handfuls of moss from Green Island [I brought this ^{vial} along to show you.]

At Cape Tuxen and more so on Green Island, the moss growth was especially luxuriant. Most of the islands, to a greater or lesser extent are "green as grass" on their northern, more or less snowfree, slopes.

Green Island, the northernmost of the

Berthelot's lives up to its advance notices in the Admiralty's Antarctic Pilot (p. 204, 2nd. edit, 1948) "... Green islet has on its northern slopes a luxuriant growth of moss nearly 4 acres in extent with peat up to 3 feet (0.9 m) in thickness. This is by far the largest unbroken patch of vegetation yet found in Antarctica." The "peat" so-called was thick, but none that I saw or walked over was quite 3 feet ^{thick} nor was the moss are so unbroken. Where the moss grew over and down the sides of a boulder or heap of rocks; it might easily give one the impression of being a moss clump or heap 3 feet thick. It was thick all right, much of it was dead or looked so underneath. There were breaks and gaps in the patch as a whole, but ^{there were} no extensive gaps or large areas on the slope bare of moss — quite a sight.

#14-63 Two dredge hauls made at ~~the~~ ^{our} Galindez Island anchorage, in 25 to 42 fms., and results combined into one. They brought up a portion of a silicious sponge, a

IV (5)

a number of worms, starfish ophiurans, a red sea-urchin and the first crinoid of the cruise, a gastropod mollusk, and a number of long stalked ascidians. A second specimen of the same species of crinoid was found in mud attached to the anchor when it was raised on 31 January.

The fish traps disappointed us, no fish, nemertean worms, a few, and a starfish - the ~~cove~~^{cove} but the Base landing was evidently a poor location for ~~fish~~^{catching}. Had there been time a second set ~~might~~^{in another location} have brought us better luck. We had drawn a blank as regards fish in Arthur Harbor the first time around. It was the second at Arthur Harbor that turned up the best ~~best~~ catch of the cruise.

This area with its many islands and its nearby "deeps" and Channels (Lemaire) is undoubtedly, biologically a very rich one. For a possible satellite station in this area a site with a good boat shelter should be selected, Winter Island - Stella Creek

Where the Penola wintered, Petermann Island where Charcot holed up in 19 , or possibly Cape Tuxen at which another look might be had for Faunal and floral growths — it is hard by Green Island. But then again, I do think we should leave the area largely or wholly to the British; They were here first, have done a lot, and will undoubtedly do more as time goes on. Not long before our arrival they the folks at the Base had sent off quite a lot of "pickled" invertebrates to the British Museum. Here should be recorded also, that to help me out with my shortage of bottles ~~which~~ they gave me a case of "handsome" wide mouth gallon jars — a "God-send", if ever a biologist had one!



February 1-3, ^{VI}①
March 2, 1963

Paradise Harbor area — including the
Chilean Gabriel Gonzales Videla Base and
the Argentine Almirante Brown station

This area is said to have breeding colonies
of three species of penguins, I have no rea-
son to doubt this but I saw only two,
Gentoo and Chinstrap, ~~and~~ in goodly
numbers and in several places. Members
of the Chilean Base said the Adellies are
not far away at "Paterna" which was
written out for me. This locality, if it is
one, I have not been able to locate. I
expect by correspondence later to run
it down. There are extensive shag (blue-eyed cormorant) colonies in this
area or near-by ^{on} Cape Duthiers, "just
around the corner" on the way to An-
xord Bay. Cape Pigeons, and about all
the birds seen elsewhere. <

Insert
here
p. 1a

Mosses collected yielded Collembola
in modest quantity.

#17-63

#24-63

Two dredge hauls made on rocky bot-
tom, 33 + 41 fms. returned a variety of
invertebrates, perhaps the greatest variety

The Chilean station is built upon an island thickly populated by Gentoo and Chinstrap penguins, right in among them seeing what's here, what it has done to the birds, in part at least, and the filth ~~it~~ that surrounds the buildings as a result, I would recommend against locating any station on a bird rookery; it is neither fair to man or beast.

On a point across the Harbor on Le-maire Island is a penguin rookery (birds not identified - did not get close enough, and on Bryde Island or a rocky ledge or islet hard by is also a good sized penguin rookery, around the shores of which a number of seals that I take to be crab-eaters, have been hauled out.

Go back to page ①
for continuation

seen so far: Sponges, hydroids, and another type of coelenterate polyp that I cannot at the moment readily place, bryozoans (~~or~~ moss animals), ophiurans holothurians, clams, 2 caprellid amphipods — elusive creatures — the only two recovered from any of our dredgings, and ascidians again. Palmerland waters if not the rest of Antarctic is certainly sea-squirt heaven. I have collected many over the years but never so many different kinds, so structurally different (external appearance) on any ^{previous} ~~one~~ cruise.

#20-63
#25-63

Of two sets of the fish traps, ^{the} first off the Chilean Base we got nothing but a "mess", a 100 or more nemertean worms and several different species of amphipods. A large jelly fish and a ctenophore "Comb-jelly" were pick-up drifting by in the tidal current.

With the dip net a lot of algae and amphipods, a large red species (Paradexamine?) were captured. These

could well furnish ample meals for many penguins. I get the idea from the penguin behaviour here. As we were working along this stretch of shore several penguins were seen actively diving, duck fashion, heads down, tail up, and when their heads came up above water again, working their bills as though they were eating something that either tasted good, or was satisfying. After a number of such dips, these birds, 3 or 4 of them headed out to sea. I did want to have a look ^{into} ~~at~~ those particular penguin stomachs to verify my suspicion that this omnipresent and abundant food material formed an important part of the penguin dietary. If the ~~sea~~ amphipods are as abundant elsewhere as in the red seaweeds here, they must play a far more important role in the economy of penguin life than heretofore realized (or has this side of the penguin dietary already been studied by some one else).

As far as our downnetting efforts are concerned "krill", the Euphausian "shrimp", have been elusive animals, but as luck would have it a goodly number were thrown up on the fore-castle deck in the small hours of the morning on our way to Paradise Harbor; one was still alive and swimming about in the depression ^{under} ~~the~~ a flush hatch-handle. From the anchor the same morning a small, white holothurian.

I am much taken with Paradise Harbor and the opportunities that it may have for biological research. I ~~feel~~ ^{am} sure that biologically at least it will justify its name.

The harbor has its ice conditions but I read in the coast pilot or sailing directions, that though there is considerable movement of ice through the passage between Lemaire and Bryde Islands, the upwards of 3 knot tidal currents

prevent ice in the Bay from freezing even in the winter months (Could be). Bergs there are, and more will be calved but do not let us forego the studies that should be made of animals under natural conditions just because of some floating ice. We should have the whole year round picture so far as possible, not just a series of summer sketches.

Moreover Paradise Harbor indents the mainland, not one of its many off lying islands.

There is an excellent station site that of the Argentine Almirante Brown Base, with not unworthy boat shelter nearby. Hope we can get title to the place and erect entirely new buildings. Against the high cliff face, I should like to see a multi-story building 2 or 3 if not more stories, with a cantilever platform "out front" to which the

The station boat could be hoisted, as well as supplies that could be stored right in the first story of the building.

Such a building as with the cliff to which I would attach or anchor it would be as snow free as the cliff. All problems of waste and water would be solved (snow and ice for melting are atop the cliff or close by - plenty of ice & bergy-bits float by. The several storied building would make for conservation of much of the warmth (heat) lost through the roof of a one story building. Then again the side of the building not being exposed to the winds would also minimize heat loss.

Danco — Cuverville Island area:

Here is an outstanding bird area —
Gentoo and Chinstrap penguins in large
rookeries. The former are centered
~~on Cuverville~~ ^{on the} and thus negate this
excellent building site as a place
for a laboratory. There also happens
to be a fine safe boat shelter on the
east side of this island behind a
natural breakwater upon which a
small shelter could be erected. At
high tide it would be isolated as
water flows in over the ⁱⁿ⁻shore end of
this ridge. Atop its widest part is
an abandoned whale boat. The whalers
knew a good shelter when they saw one.
There is quite a bit of snow free rock
here. Jack Crowell has walked over it.

Insert from
page 1a

Last but not least is the Chin strap pen-
guin rookery or rather metropolis on
Cape Spigot. It is a sight worth
travelling to Antarctica to see. On
the Argentine charts the name is Nuna-

A short distance to the north of the
Base "O" hut, ^{on Dingo Island} there is a large Gentoo
colony, ~~with~~ ^{where} birds ^{were} marching
back and forth in deep "ruts" or tracks
~~the~~ all the while we were ashore. The
landing is poor here because ~~the~~ a-
long the shore the water is so very
shallow for quite a distance out.

[Go back to page 1]

VI
(2)

Tak Negro. Its fire shore is crowded with Chinstraps, among whom a single Bendo was spotted, but the bulk of the Chinstraps in this vast rookery had nests, or roosts if you will, all over the rock, steep slopes of Cape Spigot. Every snow free patch of rock, and the greater part of this great and impressive Cape was crowded with penguins. Capt ~~Mac~~ MacDonald estimated that this ~~na~~ nunatak must have at least a 100,000 Chinstraps. What hikers and climbers the Chinstraps must be to avail themselves of this high and steep sided peak to colonize its uttermost heights 938 feet above the sea! Most of the road upwards for those that had not settled on the fire shore was over a small steep to snow field lying ~~the~~ shoreward of the bare crest of the "saddle" between the peak of the Cape and the ice sheet or cap farther to the north of it.

#27-63 Two dredge hauls were made on
#28-63 February 5; The first at the ~~anchorage~~ ^{anchorage} off
Base O, Panto Island in 41 fms, the second
at the anchorage in Errera Channel
nearer Couvreville in 46 fms. At both
the bottom was more or less rocky and
very much alike in their sampling of the
channel bottom animal life: Echino-
derms predominated, many ophiurans
a few starfish, and a dozen or more
red sea-urchins. In the second haul
were several forms of animal life not
found in the first, a large nereid worm
a number of bryozoan fragments, and
what made me want to let out a
cheer a hippolytid shrimp the first
and as it proved to be later the first
and the only decapod crustacean taken
on this cruise. Two white starfish came
up on the anchor at the first of our
anchorages of this day. A small silver
sided fish, also unique in our collections,
6 inches long, apparently dropped by
some bird, was picked up on Couvreville
while he was ashore surveying a pos-

sible site. No really suitable site for a station of any size seems available either here or at Cape Spigot unless one wants to bed down with penguins. The possibility of finding space enough for some small shelter on the ^{natural} break-water on the east side of Couvreville was mentioned above. The only need would be for penguin studies, the marine fauna would be ~~research~~ a job for a vessel hereabouts, and Chin straps might be more conveniently studied in Paradise Harbor if and when one might acquire the Argentine Base there; and as we learned later at Deception Islands.

(VII) ④

February 7, 1963

On the way from Melchior to Port Lockroy the ship headed southeastward down the Schollaert Channel. On the way she entered each of the great bays indenting the northern coast ~~was~~ of Anvers Island, first Lapeyre's lying between The Dump and the Goudon Peninsula, next Patagonia Bay between the Goudon and Thompson peninsulas and lastly Fournier Bay and Inverleith Harbor which indents its eastern shore.

Some seals, not many in any case were basking on ice-cakes or small bergs in each of these bodies of water; a killer whale (orca) was seen in Patagonia Bay — little else, only a few stray birds other than the omnipresent Wilson's storm petrels, a Dominican gull or two, a Skua now and then, a tern, a giant petrel, and a snow petrel or sheath-bill. A foggy overcast somewhat

dismal day; ice cliffs round about.

Spotted no "sites" likely or unlikely.

Saw more seals than birds.

(VII)

VII (1)

February 6, 7, 1963.

Melchior Islands area, principally with reference to the Argentine station located here on Lambda Island.

There was a great deal of snow about and between some of the buildings — most of the other islands round-about and this one off the station area ^{are} heavily ice-capped. Not much chance to set up "business" here or ^{the} urge to do it so, after seeing this side. ~~There was a great deal of snow about and between some of the buildings — most of the other islands round-about and this one off the station area are heavily ice-capped. Not much chance to set up "business" here or the urge to do it so, after seeing this side.~~

No shore collecting done, no real opportunity for it while here; few birds about, some Dominican gulls, Wilson's storm petrels, and a giant petrel (brown phase.)

#32-63 Our dredge, hauled over a mud and sand bottom at 25 fms., brought us the first sponge of any appreciable size so far, a compact, siliceous one about 8 or 9 inches in diameter; also worms; a starfish; a number of sea-urchins; an alcyonarian; a bright yellow sea-slug; half a dozen sea-squirts; and a number

of ~~tiny~~ red amphipods that either lived on ~~the~~ ^{that siliceous sponges} rough surface or in its canals.

#33-63 From the fish traps, after an over night soak in 7-11 fms., 22 fish ranging from 11 to 14 1/2 inches were found, together with 2 starfish and a "crop" of ^{the} omnipresent amphipods. Four other fish were taken over the ship's-side on hand lines. On the anchor, as it was hoisted were two bright yellow, prickly (soft-spined) ascidians like the bright yellow sea slug in color, also the first of their kind seen on this trip. Is there any reason for this identical color in two widely different forms of animal life — from the same general area, and at that not so far apart.

Lambda was the only island visited during the short time spent in this area. From a marine biological point of view the biological potential is high and holds promise of even more interesting things to come than bright ^{yellow} ~~yellow~~ invertebrates representing two widely different phyla.

VIII

February 8-10^{VIII ①}
1963

Wilhemina Bay - Svend Foyn Harbor -
~~Magdalen Bay~~ (Salvensen Cove area:

In the afternoon of the eighth, the ship moved into Wilhemina Bay past Cape Anna. The Cape is reported to be the nesting site of countless cormorants and gulls but distance and not the best visibility precluded identification of the birds whose rookeries we glimpsed.

Capt. McDonald who flew a helo recon. of Wilhemina Bay while the ship was lying-to reported: Ice cliffs on all sides, no building sites or boat shelters, 2 active glaciers, some fast ice.

The night of February 8 the ship anchored in Svend Foyn Harbor at 19:40. We then took out our fish traps and made a downnet haul. The dredge came up empty this evening, so the haul was not counted and no other attempted because of the lateness of the hour.

91 Shore collecting ^{also} proved impractical this evening and the next day ~~when~~ when a successful dredge haul was completed and the fish traps lifted. ^{over a/sand and gravel bottom}

#37-63 Dragged from 18 to 25 fathoms, the dredge bucket brought in a foot long nemertean worm; several annelids; some small white starfish; a considerable number of red sea-urchins of the species we have taken on a number of occasions; a rather large ophiuran differing, so far as I could make out, from all previously caught; a number of bryozoan fragments; the first brachiopods of the cruise, quite tiny fellows though; 2 species of mollusks; a few amphipods (as usual); and about a dozen sea squirts.

#36-63 Thirteen notothoniid fish were trapped. ~~Of~~ these the largest measured 15 inches in length, the smallest, of apparently another species 5 1/4. A 10 inch notothoniid was caught over the ship's side on a hand-line.

No shore collecting

February 9, Comdr. Lewis gave me a photo-copy of this area, on which he had noted the penguin colonies he had seen in the course of two days of helo recon flyings; he also noted ~~my~~ conspicuous lichen stands that caught his eye - encrusting yellow lichens primarily. The flights ranged from Nansen Island northward as far as

~~the~~ Challenger Island just above Bluff Island. ~~His observations were made here~~

Over ~~with the following observations:~~

- ① North end of Delaite Island, small penguin colony.
- ② Two rocks or islets, south of the largest of the Racovitza Islands have each small penguin colonies on them.
- ③ [On the way into the cove in Svend Foyn Harbor in which the wreck lies, a rocky point to the left has a shag colony on it - half the birds were sitting perhaps better standing in the sheet of snow above the bare rock area on which the rest of the birds stood. (This ~~is~~ ^{is} my personal observation ~~entered on~~ ^{entered on} the Commodore's charts.)

His observations as entered on
the charts herewith follow. He
considered as "small" colonies esti-
mated to contain under a 1000
birds, and as "medium" if the
individuals present were estimated
~~to be~~ to number from ~~an~~
1000 to 5000.

- ④ Penguins on rocks and islets ringing Icarus Point
- ⑤ and at the western end of Bancroft Bay numerous shag colonies
- ⑥ left "arm" of Reclus Peninsula on a 500 foot high cliff (west of the 1270 elevation on the chart) a medium sized penguin colony.
- ⑦ Southernmost of the Gaston Islands medium sized penguin colony.
- ⑧ Shags and sparse lichens ^{growth} on Andree Island in Recess Cove.
- ⑨ on peninsula extending ^{out from shore before} Santos Peak a small penguin colony either side.
- ⑩ northern extremity of Bluff Island small penguin colony, and lichens.
- ⑪ on Challenger Island, many lichens.

The Commadore also indicated that there were many seals on, or about the rocks (and/or ice) and islets north of the Reclus Peninsula, and ^{about} the little hooked peninsula extending eastward ~~toward the~~ Portal Point, and more seals a little farther south along the coast.

Svend Foyn Harbor was much favored by whalers in by-gone days. It is said to be a place of generally favorable weather (H.O. Sailing Directions, p. 194), even though there are at times violent south easterly winds. After having seen all three places ^{in this area} Capt. McDonald believes that ice conditions are probably better here than at Alcock Island further up the coast, or at Welchness, Dundee Island, over on the east side of the Palmer Peninsula.

The Captain also remarked that there is a small boat shelter available here near the islet on which rest several of the old, man-powered, whale boats or what is left of them. On this small rocky islet is space that could be utilized for a small building (sub-station or shelter).

The bottom of the Harbor is rocky, and poor holding ground, a fact appreciated by those whalers for they installed moorings here and there on rocky ledges as elsewhere in places favored by them along the Palmer coast and islands.

The Staten Island moved out of Svend Foyn at 15:00 and cruised the Brabant coast the afternoon of the 9th. No site areas noted. Passed Auguste Island and Cabalesou Islet. The former is mostly snow free, the latter ^{also largely snow free} has penguins and cormorants "aboard" (fide Sailing Directions) but we were too far off to see what was what. The former small island is beset with shoals and apparently offers no shelter for small craft.

For the night we lay over in Hughes Bay and the ship's force spent the following forenoon overhauling the helicopters, effecting repairs to one of the two steam boilers, and shifting the LCVP #1's motor into the Greenland cruiser.

As the ship passed through Salvensen Cove one of the deck officers noticed several patches of "red water" but did not ~~call~~ mention the matter until some time later, too late to do anything

about it. It assuredly must have been krill which has been eluding our nets and drags except for an occasional specimen. Now I would liked to have made a tow through that "red-water."

On the starboard side going into Salvensen Bay there is quite a large cormorant colony ^{with} half the birds sitting up on the snow field above the rock exposures that marked the nesting sites. It was like the state of affairs we ^{had earlier} ~~later~~ observed on some exposed rocks at the head of a cove harboring a wreck in Svend Foyn Harbor.

Dredge haul and fish trap evidence backed up by "red water" seems to indicate that the Svend Foyn area might profitably be exploited biologically.

By 11:00 the ship was underway again, headed for Brialmont Cove

IX

February 10, 11, ^{IX}①
23 and 24, 1963

Brialmont Cove, Alcock Island, Spring Point

Several hours were spent locating a suitable and satisfactory anchorage; a considerable part of the Cove ~~is carries~~
~~as depth~~ is around 200 fms. deep.

This is quite a lively place, seals scattered about. It has been some time since we have seen quite so many ~~about our~~ ^{around any of} anchorages. They were mostly, or all crab eaters. Capt. McDonald said that he saw one leopard seal in the bay, and a number about Alcock Island ~~on which~~ ^{where} there is a large penguin colony.

We killed one of the crab-eaters for bait ~~for our fish traps~~ ^{for our fish traps}. Doty shot him, the very first shot must have severed ~~his~~ ^{the} spinal cord, ~~for the~~ ^{the} seal quivered and died. Doty's second shot was not needed but was fired as a precaution before the boat crew climbed over ~~on~~ ^{to} the ice cake where the seal was resting so that we could get this "don"; it seemed, of seal meat ^{aboard}.

~~was not a very large seal~~. In its stomach were several gallons of krill. The krill in the two ends of the stomach were distinctly different in color, and I believe due to a difference in species rather than degree of digestion. Shall check when specimens get home.

Contemplating that seal's relatively small mouth and insignificant teeth, and then the gallons of shrimp in its stomach, one is ~~almost~~ forced to conclude that the krill were so numerous and so crowded together ^{that the} ~~that~~ mass ^{of them} must have had the consistency of thick porridge. How else could that seal pick up perhaps three gallons of small shrimp in his small mouth?

Again, the usual run of birds, ~~more~~ more numerous than usual: Storm petrels, Wilson's; Cape Pigeons; Dominican gulls; ~~Sheath-bills~~ ^{Shags}; Skuas; a giant petrel; and penguins, and penguins!

The latter appear to have taken

over about all available rocks and islets about ~~the~~ ^{the snow free parts of} ~~Cove~~ and virtually ~~at~~ the whole of Alcock Island. These penguins are Chinstraps, the little fellows with the urge to climb high in this world of ours. Alcock's difficult and precipitous terrain reminds one of Cape Spigot in the Danco ~~Island~~ - Couvreville Island area, but nowhere nearly so high, only a bit more than 300 feet as compared with Spigott's 938. But Alcock ^{must be} ~~is~~ much more densely populated. Capt. McDonald who has reconnoitered Alcock from the air and ^{has landed on its} ~~viewed it from~~ the shores thinks ^{it} ~~is~~ has ^{with its 300,000} more penguins on it than Cape Hallett ~~down~~ McMurdo way. He said he believes it to be about the largest penguin rookery that he has ever seen in all his eight years of Antarctic experience.

At the foot of Alcock's precipitous slopes, the Captain ~~was~~ ^{afford} found a ^{for} small cove that would ~~shelter~~ ^{shelter} a

Small boat. ^{Adjacent was.} ~~and~~ a piece of penguin-free land upon which one might build without interfering with the Chinstrap way of life, which includes an "awful" lot of steep up hill climbing for they occupy most every ~~other~~ bit of ~~Alcock's~~ snow free rock on Alcock as at Spigot, to its very top.

However, I would like to see the Alcock Chinstraps, as well as those at Cape Spigot left undisturbed. Anyone having the urge to study ^{Chinstraps} ~~them~~ can do so in Paradise Harbor, living as ^{in 1959-1960} ~~did~~ A.C. Bogdanovich at the Chilean station ~~there~~ ^{or even more} as an observer, or even more conveniently at Deception Island where ^{live several hundred thousand} ~~there are some~~ Chinstraps ^{where} and comfortable accommodations may be found at any one of three stations, Argentine, British or Chilean.

Mosses and lichens were collected ashore on each of our visits to the Cove, February 10-11, and 23-24, at Alcock Island and at Spring Point. In sprayed pools, or rather pockets water among the rocks on a rocky islet that is virtually a part of Alcock, ~~and for~~ ~~bag is at the bottom of the~~ we dip-netted with a fine meshed net the young stages of some amphipod, along with the algae that were growing in the same little bodies of water.

40-63 Two dredge hauls were made here.

58-63 The first on February 11, in 35 fms, off the ship where the bottom may best be described as a "regular concrete mix", sand, gravel and ~~broken rocks~~ and good sized stones. Some of the stones and larger pieces of gravel carried encrusting bryozoans. There were also a few small worms, ~~and~~ amphipods and bits of algae in the dredge. That our dredge did not bring up more

Two bridge heads were

bottom dwelling organisms may have been due to the inadequacy of our equipment for this type of bottom or that the place ~~in which~~ dredge was dropped was rather barren. Only another drag might tell. That opportunity was vouchsafed us on the 24th of February. This time there was mud mixed in with the gravel ~~at~~ ^{in a} depth of ~~18~~ 18 fathoms. The dredge when brought up was quite alive with a variety of invertebrates: 3 or more species of worms, disintegrated bits of some jelly-fish like organism; hydroids; an alcyonarian or antipatharian; both branching and latticed bryozoans; starfish; 1 tiny ophiuran; and several crinoids, the first time that we have taken them in numbers. What might

~~we have not gotten in other parts of the Cove if there had been time and the facility for carrying our oil drum dredge about ^{in it} ~~there~~ ~~the area~~ ~~with some time~~ ~~proved~~ ~~the~~ ~~wealth of~~ ~~its invertebrate life~~ ~~which~~ ~~surely is far~~ ~~greater than~~~~

We almost lost one of our fish traps here beneath a small iceberg. We were a long time getting it out from under, and when we did get it up, it was practically empty. From the two traps set we got only a few amphipods and bits of algae.

Despite the ~~poor showing~~ rather poor showing of dredge and trap, I still think that we have an area that will prove up rich if ever intensive work is carried on in it.

With a suitable vessel and the right ^{a shore based} gear, one would not need a laboratory ~~set up~~ ^{To properly work over this area I} ~~set up~~ would ~~it~~ recommend the acquisition of such a vessel.



February 12, 13
1963

Deception Island:

This place has most everything, plenty of "flat" land for building, sites and air strips, and plenty of running melt-water — even a "built-in" source of heat, and power perhaps. Thermal wells produce heat and power in New Zealand's North Island and in Italy. Why not here where ~~at low tide~~ clouds of steam arise at low tide ^{in the beach sands} down in front of the old whale factory?

For him who wishes to study Chin-straps this should be the place. Some 250,000 Chin straps live on the outer slopes of this old volcano. Excellent accommodations could probably be arranged for at any one of the three Bases or stations maintained here by as many different nations ^{Chile and the United Kingdom.} Argentina, ~~Great Britain, and~~

A number of other birds are to be found ^{nesting} here as well. Cape Pigeons are around most of the year except between August 11 and September 8. Common also is the Dominican gull, which in contrast to its fellows in the Argentine Islands much further South, is a non-resident at Deception Island. At Deception, we find also Sheath-bills (or Snow petrels) Wilson's storm petrels, the Giant petrel Swallowtailed (or Antarctic terns), and Blue-eyed shags. Gentoo penguins are occasional; sometimes "touring parties of 10 or a dozen arrive, at times accompanied by single Adelines. At times Macaroni penguins are seen; three pairs have been seen nesting among the Chinstraps. Bird-wise this is an altogether interesting place.

In the course of a brief period ashore three or four Sheath-bills were noticed picking over a pile of rocks on shore, and behaving along the water's edge like so many waders. Scrapings from the rocks were made. All that was noticed were some tiny ~~algae~~ mollusks (gastropods), and a coating of green ~~algae~~ algae.

x 3

Mosses lichens, and fresh-water algae from a melt-water stream were also gathered in; the mosses Berlesed for insects. ~~the~~

Time was not vouchsafed us for a dredge haul. It is true that with the trash and "junk" on the bottom left from the whaling days, we might have lost our dredge, but made aboard here by the Engineer's force out of an old steel oil drum, replacement would not have been difficult, and there was cable to spare on the hydrographic winch, the outer turns of which ~~had been~~ ^{had been} pretty well used already.

Fishing with the traps revealed that fishes were plentiful. Thirty-five were taken in the overnight set in 5 and 8 fms. They range in length from 11 to 21 inches; the larger fish were in the trap set in 8 fms.

Near the shore before the old wrecked whale factory and dry-works are a number of large steel tanks rusty to be sure. A couple of them have had openings cut in to them at ground level and are used as store houses. Should ever a "station"

④

or laboratory be planned for Reception, it would be an easy matter to convert one or more of the remaining tanks into laboratories and quarters. Cork insulation as we have it aboard the Staten Island, ports and doors cut in, two decks installed would give one a very substantial, comfortable already roofed building with a possible source of heat hard by the door, and water or its makings ~~also~~ close at hand. ~~⊗~~

Except for the fact that the island is already over crowded, it could be made into a very wonderful center for biological studies. The possibility that the old volcano might again act up would always be with us, but is it not a dying one, though slowly if you look back over its history?

Insert
above

There is a well, back from the beach in which the water has a temperature of 65°F. , but this is not for drinking, melt-water streams are every-where, and snow and ice ^{readily} accessible

(XI)

XI ①

February 13, 1963

Yankee Harbor:

Here there is a great deal of more or less level land upon which one could build. However, the long spit of rock, rounded boulders, stones, and gravel (shingle), strikes me as an area that can be ~~down~~ terribly wet, wind and wave swept at times. The stones ^{forming the ocean front rampart} seem to have been piled ^{up} ~~about~~ rolled about ~~and~~ as so many ^{grains of} sand ~~grain~~ in a sand dune. On land, ^{there is} no protection against storms, though the spit offers protection for boats ^{within} the harbor.

Within the harbor we found the fire-shore almost wholly blocked by drift ice, carried there, and held ~~there~~ by tidal or other currents. Though a fairly strong wind was blowing across the spit from the ocean side it had no effect on the ice sheltered from that wind by the spit low as it was. I get the impression that that ice is always

and constantly kept replenished by slides from the ice cliffs that ring the other half of the harbor.

The better building site or sites, other things being discounted so far as terrain goes, are on the penguin rookery at the head of the harbor. ~~and below the hills over~~
For this reason I recommend against this place for a permanent station.

Mosses, a few lichens, and samples of the algae on the beaches were collected. From ~~the~~ moss ~~in sections~~ scraped off of a rock high on the spit we Berlesed more mites than I had seen so far on this trip. There were some Collembola too in the mosses collected ~~from~~ ashore.

#44-63 Two dredge hauls were made at the ship's anchorage in 30 fms. The first brought ~~up a small starfish; a half dozen crinoids the greatest number of these echinoderms taken in any one place, before this we had but one or two in very few of our previous hauls; ~~the~~ ~~these~~ ~~the~~ ~~had~~ ~~was~~ ~~down~~ ~~no~~ ~~much~~ ~~no~~ ~~sand~~~~

There were also a few worm tubes; ~~the~~ ^{XL (3)}
~~the haul~~ several small ascidians and
algal fragments; All ^{were} washed clean of
bottom material, no mud, no sand, no
gravel, no rocks. In the second haul
we got a lot of stiff mud from which
~~were~~ sponges; many tube worms; ophiurans
ascidians; and algae were obtained.

Our stay ~~there~~ was too brief for a trial
with the fish trap. These should have not
less than an all-night soak; ^{besides} ~~with~~ 48
or 36 ~~hours~~ ^{hours} are always better than 24.

One is intrigued with the number of
crinoids. More drags could be profitably
made, but from a trawler, rather than
working out of a shore base ~~the~~
with a small ~~boat~~ ^{boat}.

I do not favor Yankee Harbor as
a laboratory site for reasons stated
above — penguins, wind, weather and
ice.

~~Handwritten scribbles and text, possibly crossed out.~~

XII

XII (1)
February 14, 1963

Hope Bay:

This is an icy, extremely windy place as Capt. McDonald can tell you.

There are two bases already here, British and Argentine. The latter occupies the most favorable part of the area and has the best installation, as well as landing, or dock facilities. Landing from boats at the British Base is difficult except at high-tide; a pier once there is no more.

The Argentine Base sits upon what must have been a part of the large Adelie colony nearby. Prospects are that it will, in time become further depleted as the young pups and sledge dogs (to be) still untrained, run free. While we passed through 3 young dogs ran down, worried, and killed a full grown penguin. How often this happens is anybody's guess. The Argen-

XII (2)
time who was showing us about, ~~the~~
went through the motions of pulling the
dogs off, slapped at them which
gave us the license to administer a
kick or two, all to no avail. The
dogs would turn away, ~~and~~ ^{but} ran after
the penguins ^{again} as soon as our backs were
turned. One got in a good bite — there
was no use ~~sticking~~ sticking around longer.

With untethered young dogs about
one can easily clear off a penguin
rookery for a building site. I do
not recommend this, nor write this
for publication — this paragraph and
the one preceding it!

At the Argentine Base the Wardroom
mess was hospitably entertained at
an "Asado", a real "criollo" affair.

At the British Base we had earlier
had tea. Here we learned that oc-
casionally Chinstrap ^{and a few Gentoos} turn up; that
Sheath-bills are resident, Skuas and
Dominican gulls, well known; Snow
petrels are occasional, as are also

Fulmers and Giant petrels; shags occur in limited numbers, and Antarctic terns (or are they the Swallow tailed ones) can be found about the "lakes". Young's Point has Silver-grey petrels, and Adelie's have a rookery on ~~some~~ Island further south; still further on one encounters Emperors.

High tide curtailed our dipnetting along shore. Moss was about in very small patches, widely scattered; did not see any worthwhile "stands". What we gathered and Berlesed yielded very few insects.

The dredge haul at the ship's anchorage, in 25-30 fms, mud and sand bottom, turned up many tube worms, some small clams, a gastropod, ophiurans and algae.

Twenty-two fish were trapped, a goodly catch — where there are fish in plenty there must be an abundance of food for them. Stomach contents of those picked out for preservation,

I expect to have examined in Washington. Having no facilities for curing for a lot of pickled fish, the fish I have saved have been frozen. Let us hope that we can get them all back in the same state.

I cannot recommend Apo Bay as a possible or future station site. I am sure it is located in a biologically rich and interesting area, but as already said it has, ~~been pretty well~~ been taken over by others. On the score of wind and weather too, I cannot rate it high in any list of station sites.

February 15, 17-19,
1963

Seymour Island - Snow Hill area:

Seymour Island has about as large or perhaps larger snow free area as any place we have seen to date. It is ~~not~~ a soft and muddy place; one could at most call the soil, loam, so soft is it, but nothing much grows on it. ~~I~~ ^Saw and gathered very little plant material ashore.

"Amid-ships" toward the South is a great "alluvial" river valley with running water and tributary streams, some though at this time of year ~~were~~ much reduced in volume.

Few birds, fewer seals, about ~~to~~ ^{one} dead one, very old carcass, only skin and bones left, was found in a gully toward south end of the island.

The island is ~~quite~~ a barren looking place ~~so far as animal and plant life was concerned when we were there.~~ so far as animal and plant life was concerned when we were there.

Paleontologically the island has quite a different "face". Much has been done here in this line. I do not know how much more remains to be done. This would bear looking into. Mr. Berg, and the Commadore ~~in~~ a brief space of time secured a number of highly interesting fossils, wood, bone, and shell specimens.

Most zoologists are interested in paleontological doings, for them it is but fossil zoology.

As to recent organisms: From the ship lying to in the ice off Seymour Island a few hours on Feb. 15, two hauls were made in 38 fms over a rock, sand, and gravel bottom, because they were made at the same place and were so similar, the results were treated as one. Thin thread-like worm tubes were in this drag by the thousands, along with some clam, and nemertean worms; included in the haul was our first living barnacle; some alcyonarians; and several stalked, white

ascidians that had much the look of "Jack-in-the-pulpits".

Another dredge haul ~~was~~ made while at anchor off Snow Hill Island on the 17 Feb. in 14-15 fms., on a bottom of mud so tough and stiff that it was like handling silicon putty, or ^a tough synthetic rubber mass.

In this haul we found a strange polyp, the type of coelenterate represented I do not know; there were also a nemertean worm, 3 species of annelids; several sea-urchins in part fragmented; some bivalve mollusks, and the first cumacean so far seen. (Cumaceans are small, often tiny shrimplike crustaceans, bottom feeders.)

Two other dredge hauls combined, ~~at~~ were made in 12 fms over a tenacious mud-cemented sand bottom on the 19th. This time we got more worms; a starfish; hydroids; a piece of an alcyonarian; one tiny amphipod; and a number of small living clams, and some dead shells.

Perhaps the most interesting of all specimens that were secured in this

area were three large red sea-spiders Pycnogonids. These had hopped onto a fish trap let down over the ^{side of the} anchored ship on the night of the 18th, supposedly a fathom or two off the bottom so that the trap would not drag as the ship swung with wind and/or tide. The trap hauled up the morning of the 19th ~~contained~~ ^{held} nothing else of ~~much~~ consequence, a fragment of a yellow sponge, and scraps of algae.

Tried this "stunt" the night of the 19th but lost the trap as some ice-cake or berg coming alongside during the night carried it off. There seems to be quite a bit of drift ice in desultory movement in this area, ~~probably~~ ^{probably} more wind and current driven than otherwise, but its behaviour could not be foretold or counted on.

We really have a rich and promising bottom in this area, in and about Seymour, Snow Hill, and ^{islands} ~~nearby~~ Vega, Cockburn and others. In almost every dredge haul made along the Palmer Penin-

XIII (5)

sula we got some form of animal life that we had not turned up in any of the hauls preceding the one in hand.

It does seem to me that a thorough-going ^{marine} biological survey be made of the territory the sea and bay bottoms sailed over in this cruise, in the areas where the El Panin cannot or should not operate.

And for this a vessel or vessels should be provided with all necessary "gear" and equipment enabling it to work in depths upwards of 400 and 500 fathoms and with auxiliary small craft for inshore, shallow water work.

(aboard)

(XIV)

XIV (1)

February 20, 21,
1963

Welchness, Dundee Island:

Here we have acres of snow free land, enough or almost enough to set-up a second McMurdo, but the place apparently has been taken over by the Argentines judging from buildings, survey stakes, and tractor tracks.

Landing with or from the LCVP was troublesome, beached ice-cakes lined the shores of the peninsula, or spit that forms the major part of Cape Welchness, ~~landward~~ ^{leeward} as well as windward side.

41

Ashore we covered the open, more or less level, snow-free land from one side of the peninsula to the other. Found only scanty and scattered ^{small} patches of moss. Apparently few birds about, other than a small flock or group of terns where the peninsula met the ice cliffs on the ~~seaward~~ ^{windward} side, none of the birds seen were nesting; Gentoo penguins were few and far between; a Dominican gull or two was flying

more were "roosting" on an ice cake ^{just} off shore ^{XIV} ②
about, ~~and~~ Standing on a small hillock
~~was a~~ Skua, and ^{her} nearly mature
chick, but no others in the air at the
time. The garbage dumped ~~from~~ from
the ship always brought more birds
around than we saw on, or over land,
excepting of course penguins.

Some live seals, not many were lying
on the beaches, more ^{were} farther out on
ice-cakes, but ~~farther~~ ^{well} inland, and
scattered far and wide over the whole
area were 20 or 30 (perhaps more)
remains of dead seals. I would judge
adolescent or half grown — may have
been younger. ^{the} Skin, and bones with-
in ^{it} ~~were~~ were about all that was left; heads
seemed ~~to~~ do have been picked clean. The
cause of death — I find it difficult to
believe that they ^{could have been} ~~were~~ still born young —
were much too large.

#53-63 A forenoon and afternoon dredge
haul were combined and treated as
one; were made in 30 fms at the
ship's anchorage, mud bottom. As on
such bottoms, worm tubes formed the

bulk of the catch; a half dozen species of annelids were represented; one small sipunculid worm; stardishes, ophiurans, and the largest holothurian yet taken; there were also a few coelenterate animals, stalked ones attached to small pieces of rock; and two long rope-like strands of a colonial ascidian. Another of ~~these~~, $1\frac{1}{4}$ inch in diameter "ropes" was brought to me by a member of the ship's recreation party that was put ashore in the late afternoon. Also a dried pycnogonid (Decapoda antarctica) was picked up on the beach; as well as a specimen of "krill" from near the mouth of a small melt-water stream flooded by the tide. Of the great windrows of algae on the beach a few samples were saved.

Here we lost the two traps that we had regularly been using of late. So many and so large ice bergs moved in on shore that we could not locate the ^{trap} floats. ~~we still had~~ ~~the~~ ~~one~~ one in reserve ~~and~~

The engineering department on the Staten Island quickly constructed us another coil of wire that I had purchased in New Zealand for just ~~that~~ such a contingency. ^{But} ~~undertake~~ there was no opportunity to ~~do~~ a second trial here. ~~with fish traps.~~

Though the biological potential here—abouts from the marine biological point of view, rates good or better, bird and plant life are scanty. ⊗

There are fossils to be found. ~~and~~ We ~~did~~ got a fine specimen of an ancient clam. Unless the paleontology ^{covered} ~~has been~~ by others, there ^{here} ~~still~~ may be an opportunity to accomplish something in this line at Welchness.

The ^{apparent} pre-occupation of the site, by Argentina, however, does argue against ~~a~~ a U. S. biological station here at this time.

* I think Landon report remarks that some marine biological work was carried on here in 1951-52. Shall check.

(XV)

February 25^①
1963

False Bay, Livingston Island:

Weather none too good, had to wait it out before we could be landed ashore even by helicopter; no place in this weather to beach shore boat; surf or swell breaking on the steep-to shingle and boulder covered beach (moraine debris?) rendered landing by boat impossible! — at least in this, and I would say ⁱⁿ much other weather either. Getting in and out, off and on shore is a must for any station, and this should be possible most of the time. If not, the place is out as a station site.

#60-63 Did gather a fair collection of lichens and moss; Berlesed the latter aboard. Not time enough ^{spent} here to warrant setting fish traps.

Chinstraps were the only penguins seen ashore a couple of rather widely ^{of a dozen or so} separated lonesome little groups sitting forlornly among the rocks. No evidence

XXV (2)

that there had been a rookery on this beach. These penguins were probably ^{of} this year's crop "on their own" having been cast out at home; some of them were half through moulting their "chick" plumage.

Around the "corner" in South Bay, Commodore Lewis spotted a number of seals resting as usual, but this time in a fresh water pool. He said also that there was a colony of penguins about 400 feet up probably Chinstraps, those are the only ones we saw elsewhere in these parts — and how the little "beggars" like to climb high!

As his helicopter swooped down for a closer look, the penguins all ran toward the edge of the cliff several dived over, to be killed on the rocks below. Seeing this

#61-63 In the dredge: muddy coarse sand, rocks and gravel from 17 fms. along with hydroids; no end of worm tubes of various sorts; 2 ophiurans; mollusks; an amphipod; 2 isopods; 1 specimen of krill; 1 pycnogonid; some long strands of a colonial ascidian, and a few solitary stalked one.

Might enjoy dredging here; but a station in this place — No!

Commodore did not continue flying in this area because of the bad weather.

(XVI)

XVI (7)

February 25, 26
1963

Discovery Bay, - Ash Point, Greenwich Island:

Quite a wide open place, windy as "all get-out" while we were about full of shoals; weather thick to say the least.

Went out with survey party to Ash Point, hoping to be able to pick up our fish traps that had been set out the night before (February 25), but were recalled to the ship before this could be done.

Lt. Beam who went ashore with the survey party undertook to check on the penguins there. He saw no more than 5 or 6 Gentoos, and a single Chinstrap.

At this time of year, and in the weather we encountered here the Point is a bleak, sparsely, if at all populated place. In general Chinstraps seem to predominate in the rookeries so far seen in the South Shetlands, I ~~almost~~ expected to find more here. Still we did not get around as anticipated because of the weather. A number of the ship's personnel got

Feb. 25

stranded ashore, and an equal number of Chileans spent the night aboard the Staten Island when the seas and wind got too high the evening of the **25th**

#62-63

on the 26th

A dredge haul in 31 fms, mud bottom, at ~~the~~ ship's anchorage in the Bay turned up the first sizeable brachiopod of the cruise. Though no larger around than a nickel, it was ever so much larger than the few smaller than pea-sized specimens we got on just one other ~~occasion~~ ^{there were} occasion; otherwise a lot of tube dwelling worms; 3-4 species of hydroids; bryozoans; a dozen extremely thin shelled snails mostly broken or crushed; ^{and} a half dozen of the rope-like colonial ascidian such as we had taken on several other occasions. This time we got the ~~whole~~ ^{complete} animals, it seems; I did not know that they had roots holding them to the bottom. Well they have — good for our old steel drum! I always thought these "fellows" were free swimming like salps.

#63-63 In the fish traps when we got them back were just 2 nototheniid fish which I took to be ~~different~~ species, both were saved, a third little orange colored fish the "boys" said ~~to have been in the trap~~ got away in the surf. fle; otherwise there was just a starfish, an ophiuran, a small, crushed, red sea urchin; and a few fragments of algae.

Except for ~~the~~ interesting sea life that seems to be ^{had} here, the area as a whole does not particularly appeal to me. Earlier in the summer there may be more animal life about, but if we are to biologically do Palmer land I would rather be closer to it.

XVII

February 28,
March 3, 4, 1963

Ardley Island area ^{and} Potters Cove,
King George Island:

This February 28, at 05:15 Capt. McDonald and I had a helicopter flight over Ardley Island and vicinity. We did not get over Potters Cove. There were lots of lakes in the hills and lowlands. Had a good view of the reef exposed at low tide that leads some to call the island, Ardley Peninsula.

~~We encountered on March 3 that~~
The bad weather ~~at yesterday that~~
ruled out all operations in Collins Harbor
continued ~~over the 5th~~ and interfered with all
plans. The survey ^{party} that went ashore,
had only 15 mins here before being
recalled.

#64-63 ~~The~~ Weather however, did not interfere
with ~~a dredge haul on the afternoon of the 28th~~ ^{dredge haul}. The very
muddy bottom over which the drag was
made, 47 fms down, must be covered,
literally with a forest of worm tubes,
the work, I should say, of some 8 to 10

different species of worms; an equal number of ascidian species were counted in his haul too, so intimately must some of them been associated with the tube worms that they had grown around the tubes, and used them for supports; there were also hydroids; 3-4 species of mollusks; 3 of sponges; 2 of starfish, and 2 of ophiurans; lastly a single isopod.

We did get our fish traps over but they were left on the bottom to be picked up on our return from Arthur Harbor where we were to go for the next day or two. It was a generally bad day when we got back ~~to them~~ on March 3.

On our way to get the traps we dropped a survey party including Mr. Crowell and Capt. McDonald on a beach between the Ardley and Fildes peninsulas.

Shortly after we got the traps up we were recalled to the ship, the idea being to give Lt. Beam a chance to

took over an old wreck somewhere in the vicinity. He did not get to go nor did we get to take off the shore party. They were left to their fate, so to speak. Notwithstanding, they had themselves a party bonfire and shelter building in anticipation of a night ashore. Only Mr. Crowell out of the entire group of 8 or 10 had any survival gear with him - no radio - no boat.

At low tide the party probably could have made it over ~~to~~ Ardley Island ~~the~~ on the connecting reef where there is a shelter hut, in case of necessity. Even so they found a food cache right where they were stranded. It had been left behind by an earlier English survey party. However no need arose to open it. For about supper time they were rescued.

The water inshore though we could not see how it was from the ship at anchor, was relatively smooth and quiet compared with the sea ~~that was bouncing~~ out where the ship was "bouncing" up and down. All's well that ends well.

XVII (4)

was Capt. McDonald who was ashore with ~~that~~ ^{the} survey party, with the "excellence" of the area ~~as a~~ ^{as a} building, for shelter for boats, and the quiet water in shore, that I believe this Fildes-Ardley Peninsula site ^{became} his second choice ^{site} for ~~setting up~~ ^{locating} the Palmer Peninsula biological station.

As this was no day for the birds and was ^{so very} late in the season, little can be said of the bird life of the vicinity. There were seals about. The question was raised, could they be for seals? There was no chance to investigate.

In the lifted traps, 6 fish were found 10 1/2 to 13 1/2 inches in length, half a dozen starfish, and a number of red amphipods.

March 4 - Potters Cove:
This forenoon, ashore ^{mosses, lichens,} clumps of grass and samples of the marine algae that had been thrown up on the beach were collected. Berlesed the mosses; later ^{out of the bag in which they had} when I took moss and grass.

placed,

XVII (5)

Few insects (flies?) dropped in the pan over which I was working, and were quickly bottled.

There were several crab-eaters hauled out on the beach. Gentoos and Chinstraps were to be seen but in very limited numbers, a few of each at most, and fledglings at that. ~~Also noted were~~ ~~between us we saw also~~ Dominican gulls, skuas, Wilson's storm petrels, and two birds that Capt. McDonald said were Antarctic petrels or perhaps ~~the~~ Giant Petrels. Lendon in his report listed Sheath-bills ^{but} we saw none.

The dredge haul of this day, in Potters Cove, in 19 fms, mud bottom, produced a fine lot of mud dwelling, tube building worms, if anything larger and "handsomer" than those we got off Ardley Island but of the same species surely, ~~most of~~ ^{likewise} the mollusks were of the same species as yesterday, but the sponges, ~~hydrozoans and~~ hydroids and echinoderms ~~present in the haul of the day before~~ we failed to find; did get an alcyonarian though.

After a morning flight over the area, Capt. McDonald remarked that Potters Cove seems to have about everything: the necessary Piedmont insuring safe landing for aircraft ^{with space} on the snow-free land too, ^{for an airstrip,} certainly for helo pads; a dock, mooring facilities, and a sizeable lake for water supply. The dredging holds forth hope for more rewarding collections ^{than} ~~than~~ we made during our brief stay here, but I, for one, would like to leave the South Shetlands, biology and geology, to the British who have already done so much in these fields in this ^{general} area, and in ^{and about} ~~these~~ ^{these} islands. ~~South Shetlands.~~

XVIII

XVIII ①

March 4, 5,
1963

Admiralty Bay:

Here too we struck bad weather, high winds and poor visibility, and were unable to effect a boat landing on either side of the Keller Peninsula. Tried the East side the afternoon of the 4th but found too much ice in the shallow water in shore. The survey party was put ashore by helicopter.

Because of continuing unfavorable winds the projected landing on the West side of the Peninsula on the 5th was also given up.

#75-63 However, a dredge haul was accomplished on the 4th. It was a rather meager one, containing a number of tube building worms, starfish; and a clam or two. The depth was 22 fms., the bottom mud.

Our fish traps put over this same evening, but not lifted until mid-morning of the 5th (having waited for the wind to die down) contained a very scanty catch, a couple of worms; some amphipods, and a starfish.

Coming up this area under very unfavorable circumstances, our trials were limited, nor could a thorough examination be made of the site on the East side of the Peninsula. However, there appeared to be land enough for an extensive installation if one had to be placed here.

The last we saw of King George Island was Penguin Island off to the East. Between poor visibility and the need for keeping a safe distance off shore in bad weather we ~~are~~^{were} not in a position to identify the bird colonies said to be on this little island and the adjacent nearby shore.

~~As~~ As we passed the spray over the ship almost obscured our sight of the island, sheets of it went right up against the wheel house windows a good 70 feet above sea level.

Thus it was as we bowed out of the Palmer Peninsula area and the South Shetlands.

5 ep. sheet
 Heading Appendix A Palmerland Survey, 1962-63 —
 Tow-net hauls made:

T-1-63	Marguerite Bay	January	18
2	Bonapart Inlet		24
3	Arthur Harbor		25
4	Port Lockroy		26
5	Petermann Island		28
6	Galindez Island		29
7	" "		31
8	Paradise Bay Alv	February	1
9	Bryde Island		2
10	Paradise Bay Alv		4
11	Danco Island		5
12	Melchior		6
13	Svend Foyn Harbor		9
14	Whalers Bay		12
15	" "		13
16	Hope Bay		14
17	Welchness		20
18	Alcock Island		23
19	Discovery Bay		26
T-20-63	False Bay		25 ⊗

⊗ was numbered wrong,
 date is correct, will do as is

Tow net hauls Palmerland Survey

T-1-63	Marguerite Bay	January	18
2	Benaparte Inlet		24
3	Arthur Harbor		25
4	Port Lockroy		26
5	Petermann Id.		28
6	Gallindez Island		29
7	" "		31
8	Paradise Bay Abr	February	1
9	Bryde Island		2
10	Paradise Abr		4
11	Danco Island		5
12	Melchior		6
13	Svend Foyn Abr		9
14	Whalers Bay		12
15	" "		13
16	Hope Bay		14
17	Welchness		20
18	Alcocks Island		23
19	Discovery Bay		26
T-20-63	False Bay		25*

* was numbered wrong,
date is correct; will do as is.

A FEW QUESTIONS AND COMMENTS
REGARDING
BIOLOGICAL WORK IN THE PALMER PENINSULA

by

Waldo L. Schmitt, Ph.D.
Research Associate
Smithsonian Institution
National Museum

Looking back over our site survey, what specifically is it that USARP hopes to accomplish with the proposed Palmerland laboratory, or "boat," either one, or both?

Are these facilities to be open, available, and unrestricted for each and every applicant desiring to engage in some phase, major or minor, of any line of research that he might fancy in the broad field of biology?

Or does USARP, to begin with, wish first to further investigations directed toward carefully considered, definitely stated objectives - researches holding forth promise, or at least the hope of yielding information being sought, or of solving specific biological problems concerned with Palmerland's inhabitants, plant or animal?

In fact, in order to determine what those facilities should consist of, does not something definitive in the way of a program or plan for the Palmerland operation need to be formulated, and made known in advance, at least for the first year or two of their existence?

Otherwise, could not a lot of effort be expended in all, and perhaps wholly unrelated directions in diverse investigations that might never contribute toward the building up of the substantial, well founded, interrelated, cohesive body of biological knowledge which should result from the research work that the National Science Foundation may be called upon to underwrite in Palmerland?

Should not "first" things come first? "The first need," as set forth for the Antarctic as a whole in the Projected Antarctic Science Program by the Chief Scientist of the Office of Antarctic Programs,

Dr. A. P. Crary, "is in the field of taxonomy and systematics, the classification and description of the flora and fauna, not only of Antarctica but also of the many islands of the high southern latitudes and even the southern mainlands of other continents. Along with this description should go ecologic studies of the environmental and climatic factors under which the flora and fauna live...."

Do we agree that these things should come first in any program set up for Palmerland biological studies?

If so, is not a comprehensive, thoroughgoing, biological survey of an area representative of a variety of environments and a diversity of species, in order?

Such a survey would reveal the distribution of types of bottom, the kinds of animals and plants living on them, the communities and associations they form, their abundance and seasonal occurrence, as well as the "flow" and the constitution of the waters which distribute their larval stages, and in which still other animals and plants drift about, if they are not free-swimming forms.

How best to attain these ends -- a shore station or stations out of which to operate with a small craft, thirty feet in length or less, as was attempted at the Hallet Station, I believe?

Or by a sea-going trawler type of vessel, able enough to travel back and forth from a suitable port on the South American mainland, or even the States, if need be, equipped to explore depths of from 450 to 500 fathoms, capable of working in, or at least about ice fields, and provided with laboratory space and accommodations for a scientific

staff of four, perhaps?

Such a vessel could also work the intermediate and surface waters, and by means of its auxiliaries, shallow inshore waters, and if and when necessary support "excursions" ashore for related studies, fresh water and terrestrial. This vessel may need a shore base, or supply depot conveniently located in or near the area to be surveyed.

In no way is it intended here to minimize or overlook what can be done land-wise from a shore-based laboratory. But could not microbiological studies of soil, air, and of water and studies of bird and terrestrial plant life be conducted, as well and to the same extent, from a vessel as from a shore station, if they ^{be} deemed of such vital importance that they must be initiated the very first year or two this vessel is operated?

Penguin and other ornithological researchers have the opportunity of pursuing such studies by living, as observers, in any one or another of the presently active bases in Palmerland, Argentine, Chilean, and English. The little studied Chinstrap penguins can be studied at close range, as well at Deception Island and in Paradise Harbor, as they can be in (or at) any of the colonies of these birds observed in the course of the survey cruise.

The prospect of seeing, hearing, or reading about what may come out of a biological survey of the channels, straits, bays, and harbors of Palmerland is truly a fascinating one.

- - - - -